

**APPLICATION**  
**FOR**  
**UNITED STATES LETTERS PATENT**

TO THE ASSISTANT COMMISSIONER OF PATENTS:

BE IT KNOWN THAT WE PETER B. DERVAN AND ELDON J. BAIRD

have invented certain new and useful improvements in

**"STEREOCHEMICAL CONTROL OF THE DNA BINDING AFFINITY,  
SEQUENCE SPECIFICITY, AND ORIENTATION-PREFERENCE OF CHIRAL  
HAIRPIN POLYAMIDES IN THE MINOR GROOVE**

of which the following is a specification:

**Stereochemical Control of the DNA Binding Affinity, Sequence Specificity, and  
Orientation-Preference of Chiral Hairpin Polyamides in the Minor Groove**

5 The U.S. Government has certain rights to this invention pursuant to Grant Nos.  
GM 26453, 27681, and 47530 awarded by the National Institute of Health.

**CROSS-REFERENCE TO RELATED APPLICATIONS**

10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995  
1000  
1005  
1010  
1015  
1020  
1025  
1030  
1035  
1040  
1045  
1050  
1055  
1060  
1065  
1070  
1075  
1080  
1085  
1090  
1095  
1100  
1105  
1110  
1115  
1120  
1125  
1130  
1135  
1140  
1145  
1150  
1155  
1160  
1165  
1170  
1175  
1180  
1185  
1190  
1195  
1200  
1205  
1210  
1215  
1220  
1225  
1230  
1235  
1240  
1245  
1250  
1255  
1260  
1265  
1270  
1275  
1280  
1285  
1290  
1295  
1300  
1305  
1310  
1315  
1320  
1325  
1330  
1335  
1340  
1345  
1350  
1355  
1360  
1365  
1370  
1375  
1380  
1385  
1390  
1395  
1400  
1405  
1410  
1415  
1420  
1425  
1430  
1435  
1440  
1445  
1450  
1455  
1460  
1465  
1470  
1475  
1480  
1485  
1490  
1495  
1500  
1505  
1510  
1515  
1520  
1525  
1530  
1535  
1540  
1545  
1550  
1555  
1560  
1565  
1570  
1575  
1580  
1585  
1590  
1595  
1600  
1605  
1610  
1615  
1620  
1625  
1630  
1635  
1640  
1645  
1650  
1655  
1660  
1665  
1670  
1675  
1680  
1685  
1690  
1695  
1700  
1705  
1710  
1715  
1720  
1725  
1730  
1735  
1740  
1745  
1750  
1755  
1760  
1765  
1770  
1775  
1780  
1785  
1790  
1795  
1800  
1805  
1810  
1815  
1820  
1825  
1830  
1835  
1840  
1845  
1850  
1855  
1860  
1865  
1870  
1875  
1880  
1885  
1890  
1895  
1900  
1905  
1910  
1915  
1920  
1925  
1930  
1935  
1940  
1945  
1950  
1955  
1960  
1965  
1970  
1975  
1980  
1985  
1990  
1995  
2000  
2005  
2010  
2015  
2020  
2025  
2030  
2035  
2040  
2045  
2050  
2055  
2060  
2065  
2070  
2075  
2080  
2085  
2090  
2095  
2100  
2105  
2110  
2115  
2120  
2125  
2130  
2135  
2140  
2145  
2150  
2155  
2160  
2165  
2170  
2175  
2180  
2185  
2190  
2195  
2200  
2205  
2210  
2215  
2220  
2225  
2230  
2235  
2240  
2245  
2250  
2255  
2260  
2265  
2270  
2275  
2280  
2285  
2290  
2295  
2300  
2305  
2310  
2315  
2320  
2325  
2330  
2335  
2340  
2345  
2350  
2355  
2360  
2365  
2370  
2375  
2380  
2385  
2390  
2395  
2400  
2405  
2410  
2415  
2420  
2425  
2430  
2435  
2440  
2445  
2450  
2455  
2460  
2465  
2470  
2475  
2480  
2485  
2490  
2495  
2500  
2505  
2510  
2515  
2520  
2525  
2530  
2535  
2540  
2545  
2550  
2555  
2560  
2565  
2570  
2575  
2580  
2585  
2590  
2595  
2600  
2605  
2610  
2615  
2620  
2625  
2630  
2635  
2640  
2645  
2650  
2655  
2660  
2665  
2670  
2675  
2680  
2685  
2690  
2695  
2700  
2705  
2710  
2715  
2720  
2725  
2730  
2735  
2740  
2745  
2750  
2755  
2760  
2765  
2770  
2775  
2780  
2785  
2790  
2795  
2800  
2805  
2810  
2815  
2820  
2825  
2830  
2835  
2840  
2845  
2850  
2855  
2860  
2865  
2870  
2875  
2880  
2885  
2890  
2895  
2900  
2905  
2910  
2915  
2920  
2925  
2930  
2935  
2940  
2945  
2950  
2955  
2960  
2965  
2970  
2975  
2980  
2985  
2990  
2995  
3000  
3005  
3010  
3015  
3020  
3025  
3030  
3035  
3040  
3045  
3050  
3055  
3060  
3065  
3070  
3075  
3080  
3085  
3090  
3095  
3100  
3105  
3110  
3115  
3120  
3125  
3130  
3135  
3140  
3145  
3150  
3155  
3160  
3165  
3170  
3175  
3180  
3185  
3190  
3195  
3200  
3205  
3210  
3215  
3220  
3225  
3230  
3235  
3240  
3245  
3250  
3255  
3260  
3265  
3270  
3275  
3280  
3285  
3290  
3295  
3300  
3305  
3310  
3315  
3320  
3325  
3330  
3335  
3340  
3345  
3350  
3355  
3360  
3365  
3370  
3375  
3380  
3385  
3390  
3395  
3400  
3405  
3410  
3415  
3420  
3425  
3430  
3435  
3440  
3445  
3450  
3455  
3460  
3465  
3470  
3475  
3480  
3485  
3490  
3495  
3500  
3505  
3510  
3515  
3520  
3525  
3530  
3535  
3540  
3545  
3550  
3555  
3560  
3565  
3570  
3575  
3580  
3585  
3590  
3595  
3600  
3605  
3610  
3615  
3620  
3625  
3630  
3635  
3640  
3645  
3650  
3655  
3660  
3665  
3670  
3675  
3680  
3685  
3690  
3695  
3700  
3705  
3710  
3715  
3720  
3725  
3730  
3735  
3740  
3745  
3750  
3755  
3760  
3765  
3770  
3775  
3780  
3785  
3790  
3795  
3800  
3805  
3810  
3815  
3820  
3825  
3830  
3835  
3840  
3845  
3850  
3855  
3860  
3865  
3870  
3875  
3880  
3885  
3890  
3895  
3900  
3905  
3910  
3915  
3920  
3925  
3930  
3935  
3940  
3945  
3950  
3955  
3960  
3965  
3970  
3975  
3980  
3985  
3990  
3995  
4000  
4005  
4010  
4015  
4020  
4025  
4030  
4035  
4040  
4045  
4050  
4055  
4060  
4065  
4070  
4075  
4080  
4085  
4090  
4095  
4100  
4105  
4110  
4115  
4120  
4125  
4130  
4135  
4140  
4145  
4150  
4155  
4160  
4165  
4170  
4175  
4180  
4185  
4190  
4195  
4200  
4205  
4210  
4215  
4220  
4225  
4230  
4235  
4240  
4245  
4250  
4255  
4260  
4265  
4270  
4275  
4280  
4285  
4290  
4295  
4300  
4305  
4310  
4315  
4320  
4325  
4330  
4335  
4340  
4345  
4350  
4355  
4360  
4365  
4370  
4375  
4380  
4385  
4390  
4395  
4400  
4405  
4410  
4415  
4420  
4425  
4430  
4435  
4440  
4445  
4450  
4455  
4460  
4465  
4470  
4475  
4480  
4485  
4490  
4495  
4500  
4505  
4510  
4515  
4520  
4525  
4530  
4535  
4540  
4545  
4550  
4555  
4560  
4565  
4570  
4575  
4580  
4585  
4590  
4595  
4600  
4605  
4610  
4615  
4620  
4625  
4630  
4635  
4640  
4645  
4650  
4655  
4660  
4665  
4670  
4675  
4680  
4685  
4690  
4695  
4700  
4705  
4710  
4715  
4720  
4725  
4730  
4735  
4740  
4745  
4750  
4755  
4760  
4765  
4770  
4775  
4780  
4785  
4790  
4795  
4800  
4805  
4810  
4815  
4820  
4825  
4830  
4835  
4840  
4845  
4850  
4855  
4860  
4865  
4870  
4875  
4880  
4885  
4890  
4895  
4900  
4905  
4910  
4915  
4920  
4925  
4930  
4935  
4940  
4945  
4950  
4955  
4960  
4965  
4970  
4975  
4980  
4985  
4990  
4995  
5000  
5005  
5010  
5015  
5020  
5025  
5030  
5035  
5040  
5045  
5050  
5055  
5060  
5065  
5070  
5075  
5080  
5085  
5090  
5095  
5100  
5105  
5110  
5115  
5120  
5125  
5130  
5135  
5140  
5145  
5150  
5155  
5160  
5165  
5170  
5175  
5180  
5185  
5190  
5195  
5200  
5205  
5210  
5215  
5220  
5225  
5230  
5235  
5240  
5245  
5250  
5255  
5260  
5265  
5270  
5275  
5280  
5285  
5290  
5295  
5300  
5305  
5310  
5315  
5320  
5325  
5330  
5335  
5340  
5345  
5350  
5355  
5360  
5365  
5370  
5375  
5380  
5385  
5390  
5395  
5400  
5405  
5410  
5415  
5420  
5425  
5430  
5435  
5440  
5445  
5450  
5455  
5460  
5465  
5470  
5475  
5480  
5485  
5490  
5495  
5500  
5505  
5510  
5515  
5520  
5525  
5530  
5535  
5540  
5545  
5550  
5555  
5560  
5565  
5570  
5575  
5580  
5585  
5590  
5595  
5600  
5605  
5610  
5615  
5620  
5625  
5630  
5635  
5640  
5645  
5650  
5655  
5660  
5665  
5670  
5675  
5680  
5685  
5690  
5695  
5700  
5705  
5710  
5715  
5720  
5725  
5730  
5735  
5740  
5745  
5750  
5755  
5760  
5765  
5770  
5775  
5780  
5785  
5790  
5795  
5800  
5805  
5810  
5815  
5820  
5825  
5830  
5835  
5840  
5845  
5850  
5855  
5860  
5865  
5870  
5875  
5880  
5885  
5890  
5895  
5900  
5905  
5910  
5915  
5920  
5925  
5930  
5935  
5940  
5945  
5950  
5955  
5960  
5965  
5970  
5975  
5980  
5985  
5990  
5995  
6000  
6005  
6010  
6015  
6020  
6025  
6030  
6035  
6040  
6045  
6050  
6055  
6060  
6065  
6070  
6075  
6080  
6085  
6090  
6095  
6100  
6105  
6110  
6115  
6120  
6125  
6130  
6135  
6140  
6145  
6150  
6155  
6160  
6165  
6170  
6175  
6180  
6185  
6190  
6195  
6200  
6205  
6210  
6215  
6220  
6225  
6230  
6235  
6240  
6245  
6250  
6255  
6260  
6265  
6270  
6275  
6280  
6285  
6290  
6295  
6300  
6305  
6310  
6315  
6320  
6325  
6330  
6335  
6340  
6345  
6350  
6355  
6360  
6365  
6370  
6375  
6380  
6385  
6390  
6395  
6400  
6405  
6410  
6415  
6420  
6425  
6430  
6435  
6440  
6445  
6450  
6455  
6460  
6465  
6470  
6475  
6480  
6485  
6490  
6495  
6500  
6505  
6510  
6515  
6520  
6525  
6530  
6535  
6540  
6545  
6550  
6555  
6560  
6565  
6570  
6575  
6580  
6585  
6590  
6595  
6600  
6605  
6610  
6615  
6620  
6625  
6630  
6635  
6640  
6645  
6650  
6655  
6660  
6665  
6670  
6675  
6680  
6685  
6690  
6695  
6700  
6705  
6710  
6715  
6720  
6725  
6730  
6735  
6740  
6745  
6750  
6755  
6760  
6765  
6770  
6775  
6780  
6785  
6790  
6795  
6800  
6805  
6810  
6815  
6820  
6825  
6830  
6835  
6840  
6845  
6850  
6855  
6860  
6865  
6870  
6875  
6880  
6885  
6890  
6895  
6900  
6905  
6910  
6915  
6920  
6925  
6930  
6935  
6940  
6945  
6950  
6955  
6960  
6965  
6970  
6975  
6980  
6985  
6990  
6995  
7000  
7005  
7010  
7015  
7020  
7025  
7030  
7035  
7040  
7045  
7050  
7055  
7060  
7065  
7070  
7075  
7080  
7085  
7090  
7095  
7100  
7105  
7110  
7115  
7120  
7125  
7130  
7135  
7140  
7145  
7150  
7155  
7160  
7165  
7170  
7175  
7180  
7185  
7190  
7195  
7200  
7205  
7210  
7215  
7220  
7225  
7230  
7235  
7240  
7245  
7250  
7255  
7260  
7265  
7270  
7275  
7280  
7285  
7290  
7295  
7300  
7305  
7310  
7315  
7320  
7325  
7330  
7335  
7340  
7345  
7350  
7355  
7360  
7365  
7370  
7375  
7380  
7385  
7390  
7395  
7400  
7405  
7410  
7415  
7420  
7425  
7430  
7435  
7440  
7445  
7450  
7455  
7460  
7465  
7470  
7475  
7480  
7485  
7490  
7495  
7500  
7505  
7510  
7515  
7520  
7525  
7530  
7535  
7540  
7545  
7550  
7555  
7560  
7565  
7570  
7575  
7580  
7585  
7590  
7595  
7600  
7605  
7610  
7615  
7620  
7625  
7630  
7635  
7640  
7645  
7650  
7655  
7660  
7665  
7670  
7675  
7680  
7685  
7690  
7695  
7700  
7705  
7710  
7715  
7720  
7725  
7730  
7735  
7740  
7745  
7750  
7755  
7760  
7765  
7770  
7775  
7780  
7785  
7790  
7795  
7800  
7805  
7810  
7815  
7820  
7825  
7830  
7835  
7840  
7845  
7850  
7855  
7860  
7865  
7870  
7875  
7880  
7885  
7890  
7895  
7900  
7905  
7910  
7915  
7920  
7925  
7930  
7935  
7940  
7945  
7950  
7955  
7960  
7965  
7970  
7975  
7980  
7985  
7990  
7995  
8000  
8005  
8010  
8015  
8020  
8025  
8030  
8035  
8040  
8045  
8050  
8055  
8060  
8065  
8070  
8075  
8080  
8085  
8090  
8095  
8100  
8105  
8110  
8115  
8120  
8125  
8130  
8135  
8140  
8145  
8150  
8155  
8160  
8165  
8170  
8175  
8180  
8185  
8190  
8195  
8200  
8205  
8210  
8215  
8220  
8225  
8230  
8235  
8240  
8245  
8250  
8255  
8260  
8265  
8270  
8275  
8280  
8285  
8290  
8295  
8300  
8305  
8310  
8315  
8320  
8325  
8330  
8335  
8340  
8345  
8350  
8355  
8360  
8365  
8370  
8375  
8380  
8385  
8390  
8395  
8400  
8405  
8410  
8415  
8420  
8425  
8430  
8435  
8440  
8445  
8450  
8455  
8460  
8465  
8470  
8475  
8480  
8485  
8490  
8495  
8500  
8505  
8510  
8515  
8520  
8525  
8530  
8535  
8540  
8545  
8550  
8555  
8560  
8565  
8570  
8575  
8580  
8585  
8590  
8595  
8600  
8605  
8610  
8615  
8620  
8625  
8630  
8635  
8640  
8645  
8650  
8655  
8660  
8665  
8670  
8675  
8680  
8685  
8690  
8695  
8700  
8705  
8710  
8715  
8720  
8725  
8730  
8735  
8740  
8745  
8750  
8755  
8760  
8765  
8770  
8775  
8780  
8785  
8790  
8795  
8800  
8805  
8810  
8815  
8820  
8825  
8830  
8835  
8840  
8845  
8850  
8855  
8860  
8865  
8870  
8875  
8880  
8885  
8890  
8895  
8900  
8905  
8910  
8915  
8920  
8925  
8930  
8935  
8940  
8945  
8950  
8955  
8960  
8965  
8970  
8975  
8980  
8985  
8990  
8995  
9000  
9005  
9010  
9015  
9020  
9025  
9030  
9035  
9040  
9045  
9050  
9055  
9060  
9065  
9070  
9075  
9080  
9085  
9090  
9095  
9100  
9105  
9110  
9115  
9120  
9125  
9130  
9135  
9140  
9145  
9150  
9155  
9160  
9165  
9170  
9175  
9180  
9185  
9190  
9195  
9200  
9205  
9210  
9215  
9220  
9225  
9230  
9235  
9240  
9245  
9250  
9255  
9260  
9265  
9270  
9275  
9280  
9285  
9290  
9295  
9300  
9305  
9310  
9315  
9320  
9325  
9330  
9335  
9340  
9345  
9350  
9355  
9360  
9365  
9370  
9375  
9380  
9385  
9390  
9395  
9400  
9405  
9410  
9415  
9420  
9425  
9430  
9435  
9440  
9445  
9450  
9455  
9460  
9465  
9470  
9475  
9480  
9485  
9490  
9495  
9500  
9505  
9510  
9515  
9520  
9525  
9530  
9535  
9540  
9545  
9550  
9555  
9560  
9565  
9570  
9575  
9580  
9585  
9590  
9595  
9600  
9605  
9610  
9615  
9620  
9625  
9630  
9635  
9640  
9645  
9650  
9655  
9660  
9665  
9670  
9675  
9680  
9685  
9690  
9695  
9700  
9705  
9710  
9715  
9720  
9725  
9730  
9735  
9740  
9745  
9750  
9755  
9760  
9765  
9770  
9775  
9780  
9785  
9790  
9795  
9800  
9805  
9810  
9815  
9820  
9825  
9830  
9835  
9840  
9845  
9850  
9855  
9860  
9865  
9870  
9875  
9880  
9885  
9890  
9895  
9900  
9905  
9910  
9915  
9920  
9925  
9930  
9935  
9940  
9945  
9950  
9955  
9960  
9965  
9970  
9975  
9980  
9985  
9990  
9995  
10000  
10005  
10010  
10015  
10020  
10025  
10030  
10035  
10040  
10045  
10050  
10055  
10060  
10065  
10070  
10075  
10080  
10085  
10090  
10095  
10100  
10105  
10110  
10115  
10120  
10125  
10130  
10135  
10140  
10145  
10150  
10155  
10160  
10165  
10170  
10175  
10180  
10185  
10190  
10195  
10200  
10205  
10210  
10215  
10220  
10225  
10230  
10235  
10240  
10245  
10250  
10255  
10260  
10265  
102

occurring DNA binding proteins (Trauger, et al. *Nature* **1996**, 382, 559; Swalley, et al. *J. Am. Chem. Soc.* **1997**, 119, 6953; Turner, et al. *J. Am. Chem. Soc.* **1997**, 119, 7636). DNA recognition depends on side-by-side amino acid pairings oriented N-C with respect to the 5'-3' direction of the DNA helix in the minor groove (Wade, W. S., et al. *J. Am. Chem. Soc.* **1992**, 114, 8783; Mrksich, et al. *Proc. Natl. Acad. Sci., USA* **1992**, 89, 7586; Wade, et al. *Biochemistry* **1993**, 32, 11385; Mrksich, et al. *J. Am. Chem. Soc.* **1993**, 115, 2572; Geierstanger, et al. *Science* **1994**, 266, 646; White, et al. *J. Am. Chem. Soc.* **1997**, 119, 8756). Antiparallel pairing of imidazole (Im) opposite pyrrole (Py) recognizes a G•C base pair, while a Py/Im combination recognizes C•G.<sup>2</sup> A Py/Py pair is degenerate and recognizes either an A•T or T•A base pair (Wade, W. S., et al. *J. Am. Chem. Soc.* **1992**, 114, 8783; Mrksich, et al. *Proc. Natl. Acad. Sci., USA* **1992**, 89, 7586; Wade, et al. *Biochemistry* **1993**, 32, 11385; Mrksich, et al. *J. Am. Chem. Soc.* **1993**, 115, 2572; Geierstanger, et al. *Science* **1994**, 266, 646; White, et al. *J. Am. Chem. Soc.* **1997**, 119, 8756; Pelton, et al. *Proc. Natl. Acad. Sci., USA* **1989**, 86, 5723; Pelton, et al. *J. Am. Chem. Soc.* **1990**, 112, 1393; White, et al. *Biochemistry* **1996**, 35, 12532; Chen, et al. *J. Mol. Biol.* **1997**, 267, 1157). An Im/Im pairing is disfavored, breaking a potential degeneracy for recognition (Singh, et al. *Proc. Natl. Acad. Sci. U.S.A.* **1994**, 91, 7673; White, et al. *Chem. & Biol.* **1997**, 4, 569).

Investigators have also attempted to prevent slipped-binding motifs as well as increase DNA-binding affinity and sequence specificity by covalent linkage of polyamide subunits (Trauger, et al. *J. Am. Chem. Soc.* **1996**, 118, 6160; Geierstanger, et al. *Nature Struct. Biol.* **1996**, 3, 321; Swalley, et al. *Chem. Eur. J.* **1997**, 3, 1608; Wemmer, et al. *Curr. Opin. Struct. Biol.* **1997**, 7, 355; Mrksich, et al. *J. Am. Chem. Soc.* **1994**, 116, 3663; Dwyer, et al. *J. Am. Chem. Soc.* **1993**, 115, 9900; Chen, et al. *J. Am. Chem. Soc.* **1994**, 116, 6995). A hairpin polyamide motif with  $\gamma$ -aminobutyric acid ( $\gamma$ ) has been utilized as a turn-specific internal-guide-residue and provides a synthetically accessible method for C-N linkage of polyamide subunits (Figure 1). Head-to-tail linked

polyamides bind specifically to designated target sites with 100-fold enhanced affinity relative to unlinked subunits (Mrksich, et al. *J. Am. Chem. Soc.* **1994**, *116*, 7983; Parks, et al. *J. Am. Chem. Soc.* **1996**, *118*, 6147; Parks, et al. *J. Am. Chem. Soc.* **1996**, *118*, 6153; Trauger, et al. *Chem. & Biol.* **1996**, *3*, 369; Swalley, et al. *J. Am. Chem. Soc.* **1996**, *118*, 8198; Pilch, et al. *Proc. Natl. Acad. Sci. U.S.A.* **1996**, *93*, 8306; de Claire, et al. *J. Am. Chem. Soc.* **1997**, *119*, 7909).

Eight-ring hairpin polyamides bearing a single positively charged tertiary amine group at the C-terminus have been shown to be cell-permeable and to inhibit the transcription of specific genes in cell culture (Gottesfeld, et al. *Nature* **1997**, *387*, 202). However, recent studies of polyamide size limitations suggest that beyond five rings, the ligand curvature fails to match the pitch of the DNA helix, disrupting the hydrogen bonds and van der Waals interactions responsible for specific polyamide-DNA complex formation (Kelley, et al. *Proc. Natl. Acad. Sci. USA*, **1996**, *93*:6981; Kielkopf, et al. *Nature Struc. Biol.*, in press). Recognition of seven base pairs by ten-ring hairpin polyamids containing five contiguous ring pairings represents the upper limit in binding site sizes targetable by the hairpin motif (Turner, et al. *J. am. Chem. Soc.*, **1997**, *119*:7636). Addition of pairings of  $\beta$ -alanine with  $\beta$ -alanine, pyrrole, or imidazole has allowed extension of the hairpin motif to 8-bp recognition, as demonstrated in provisional application 60/042,222. However, those skilled in the art have recognized the extreme difficulties associated with the design of hairpin motifs recognizing longer site sizes.

The present invention involves the use of R-2,4-diaminobutyric acid as a replacement for  $\gamma$ -aminobutyric acid to make the hairpin loop. In addition, a methodology for expanding the targetable binding site size of hairpins by covalently linking existing hairpin motifs without compromising DNA-binding and sequence specificity is provided.

## SUMMARY OF THE INVENTION

This invention provides improved polyamides comprising a hairpin loop derived from  $\gamma$ -aminobutyric acid which bind to the minor groove of a promoter region of a DNA sequence. Binding of the polyamide to the DNA sequence of the promoter region inhibits expression of the requisite gene. The improvement relates to the use of R-2,4-diaminobutyric acid and derivatives of the 2-amino group to form the hairpin loop. The improved asymmetric hairpin provides for tighter binding of the polyamides to the minor groove of DNA and additionally provides an amine function for derivatizing polyamides by, for example, forming amide linkages. The improved asymmetric hairpin provides for the synthesis of tandemly-linked improved polyamides that allow for longer binding sites without compromising affinity or selectivity. The improved asymmetric hairpin may also serve to attach functional or detectable groups to the polyamide.

## BRIEF DESCRIPTION OF THE FIGURES

*Figures 1A and 1B.*  
**Figure 1A** A. Hydrogen bonding model of polyamide 1-R,  $\text{ImPyPy-(R)}^{\text{H}_2\text{N}}\text{-PyPyPy-}\beta\text{-Dp}$ , to the DNA sequence 5'-TGTTA-3'. **Figure 1B** B. Binding model of polyamide 1-S,  $\text{ImPyPy-(S)}^{\text{H}_2\text{N}}\text{-PyPyPy-}\beta\text{-Dp}$ , to the DNA sequence 5'-TGTTA-3'.

*Figure 2A and 2B.*  
**Figure 2A** Computer generated models of: (A)  $\text{ImPyPy-(R)}^{\text{H}_2\text{N}}\text{-PyPyPy-}\beta\text{-Dp}$  and (B)  $\text{ImPyPy-(S)}^{\text{H}_2\text{N}}\text{-PyPyPy-}\beta\text{-Dp}$  bound in the minor groove of double stranded DNA van der Waals surface.

*Figures 3A, 3B, 3C, 3D, and 3E.*  
**Figure 3** Structures of the 6-ring hairpin polyamides.

*Figures 4A-1 and 4A-2.*  
**Figure 4** Solid phase synthetic scheme for improved polyamides.

*Figures 5A, 5B, 5C, and 5D.*  
**Figure 5** Results of MPE•Fe(II) footprinting using improved polyamides.

**Figure 6.** Binding patterns of certain improved polyamides to a 135 bp restriction fragment.

25 **Figure 7.** Affinity cleavage experiments using improved polyamides and a 3'-<sup>32</sup>P-labeled 135 bp restriction fragment.

Figure 8. Affinity cleavage patterns of certain improved polyamides at 1  $\mu$ M concentration and 10  $\mu$ M concentrations.

Figure 9. Panels A-C represent affinity cleavage patterns of certain improved polyamides.

5 Figure 10. Quantitative DNase I footprint titration of certain improved polyamides.

Figure 11. Quantitative DNase I footprint titrations of ImPyPy-(R)  $H_2N$ - $\gamma$ -PyPyPy- $\beta$ -Dp.

Figure 12. Model for chiral hairpin folding of improved polyamides.

Figure 13. Hydrogen bonding model of a tandemly-linked polyamide.

Figure 14. Structures of exemplary twelve-ring polyamides.

Figure 15. Synthesis of tandemly-linked polyamides.

Figure 16. Quantitative DNA footprint titrations of an exemplary tandemly-linked polyamide.

Figure 17. Exemplary tandemly-linked polyamides.

15 Figure 18. Construction of plasmids pDH10, pDH11, and pDH12.

### DETAILED DESCRIPTION

Within this application, unless otherwise stated, definitions of the terms and illustration of the techniques of this application may be found in any of several well-known references such as: Sambrook, J., *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press (1989); Goeddel, D., *ed.*, *Gene Expression Technology, Methods in Enzymology*, 185, Academic Press, San Diego, CA (1991); "Guide to Protein Purification" in Deutscher, M.P., *ed.*, *Methods in Enzymology*, Academic Press, San Diego, CA (1989); Innis, *et al.*, *PCR Protocols: A Guide to Methods and Applications*, Academic Press, San Diego, CA (1990); Freshney, R.I., *Culture of Animal Cells: A Manual of Basic Technique*, 2<sup>nd</sup> Ed., Alan Liss, Inc. New York, NY (1987); Murray, E.J., *ed.*, *Gene Transfer and Expression Protocols*, pp. 109-128, The Humana Press Inc., Clifton, NJ and Lewin, B., *Genes VI*, Oxford University Press, New York (1997).

For the purposes of this application, a *promoter* is a regulatory sequence of DNA that is involved in the binding of RNA polymerase to initiate transcription of a gene. A *gene* is a segment of DNA involved in producing a peptide, polypeptide or protein, including the coding region, non-coding regions preceding ("leader") and following ("trailer") the coding region, as well as intervening non-coding sequences ("introns") between individual coding segments ("exons"). Coding refers to the representation of amino acids, start and stop signals in a three base "triplet" code. Promoters are often upstream ("5' to") the transcription initiation site of the corresponding gene. Other regulatory sequences of DNA in addition to promoters are known, including sequences involved with the binding of transcription factors, including response elements that are the DNA sequences bound by inducible factors. Enhancers comprise yet another group of regulatory sequences of DNA that can increase the utilization of promoters, and can function in either orientation (5'-3' or 3'-5') and in any location (upstream or downstream) relative to the promoter. Preferably, the regulatory sequence has a positive activity, i.e., binding of an endogeneous ligand (e.g. a transcription factor) to the regulatory sequence increases transcription, thereby resulting in increased expression of the corresponding target gene. In such a case, interference with transcription by binding a polyamide to a regulatory sequence would reduce or abolish expression of a gene.

The promoter may also include or be adjacent to a regulatory sequence known in the art as a *silencer*. A silencer sequence generally has a negative regulatory effect on expression of the gene. In such a case, expression of a gene may be increased directly by using a polyamide to prevent binding of a factor to a silencer regulatory sequence or indirectly, by using a polyamide to block transcription of a factor to a silencer regulatory sequence.

It is to be understood that the polyamides of this invention bind to double stranded DNA in a sequence specific manner. The function of a segment of DNA of a given sequence, such as 5'-TATAAA-3', depends on its position relative to other functional regions in the DNA sequence. In this case, if the sequence 5'-TATAAA-3' on the coding strand of DNA is positioned about 30 base pairs upstream of the transcription start site, the sequence forms part of the promoter region (Lewin, *Genes VI*, pp. 831-835). On the other hand, if the sequence 5'-TATAAA-3' is downstream of the transcription start

site in a coding region and in proper register with the reading frame, the sequence encodes the tyrosyl and lysyl amino acid residues (Lewin, *Genes VI*, pp. 213-215).

While not being held to one hypothesis, it is believed that the binding of the polyamides of this invention modulate gene expression by altering the binding of DNA binding proteins, such as RNA polymerase, transcription factors, TBF, TFIIB and other proteins. The effect on gene expression of polyamide binding to a segment of double stranded DNA is believed to be related to the function, e.g., promoter, of that segment of DNA.

It is to be understood by one skilled in the art that the improved polyamides of the present invention may bind to any of the above-described DNA sequences or any other sequence having a desired effect upon expression of a gene. In addition, U.S. Patent No. 5,578,444 describes numerous promoter targeting sequences from which base pair sequences for targeting an improved polyamide of the present invention may be identified.

It is generally understood by those skilled in the art that the basic structure of DNA in a living cell includes both *major* and a *minor groove*. For the purposes of describing the present invention, the *minor groove* is the narrow groove of DNA as illustrated in common molecular biology references such as Lewin, B., *Genes VI*, Oxford University Press, New York (1997).

To affect gene expression in a cell, which may include causing an increase or a decrease in gene expression, a effective quantity of one or more polyamide is contacted with the cell and internalized by the cell. The cell may be contacted *in vivo* or *in vitro*. Effective extracellular concentrations of polyamides that can modulate gene expression range from about 10 nanomolar to about 1 micromolar. Gottesfeld, J.M., *et al.*, *Nature* 387 202-205 (1997). To determine effective amounts and concentrations of polyamides *in vitro*, a suitable number of cells is plated on tissue culture plates and various quantities of one or more polyamide are added to separate wells. Gene expression following exposure to a polyamide can be monitored in the cells or medium by detecting the amount of the protein gene product present as determined by various techniques utilizing specific antibodies, including ELISA and western blot. Alternatively, gene expression following



exposure to a polyamide can be monitored by detecting the amount of messenger RNA present as determined by various techniques, including northern blot and RT-PCR.

Similarly, to determine effective amounts and concentrations of polyamides for *in vivo* administration, a sample of body tissue or fluid, such as plasma, blood, urine, cerebrospinal fluid, saliva, or biopsy of skin, muscle, liver, brain or other appropriate tissue source is analyzed. Gene expression following exposure to a polyamide can be monitored by detecting the amount of the protein gene product present as determined by various techniques utilizing specific antibodies, including ELISA and western blot. Alternatively, gene expression following exposure to a polyamide can be monitored by the detecting the amount of messenger RNA present as determined by various techniques, including northern blot and RT-PCR.

The polyamides of this invention may be formulated into diagnostic and therapeutic compositions for *in vivo* or *in vitro* use. Representative methods of formulation may be found in *Remington: The Science and Practice of Pharmacy*, 19th ed., Mack Publishing Co., Easton, PA (1995).

For *in vivo* use, the polyamides may be incorporated into a physiologically acceptable pharmaceutical composition that is administered to a patient in need of treatment or an animal for medical or research purposes. The polyamide composition comprises pharmaceutically acceptable carriers, excipients, adjuvants, stabilizers, and vehicles. The composition may be in solid, liquid, gel, or aerosol form. The polyamide composition of the present invention may be administered in various dosage forms orally, parentally, by inhalation spray, rectally, or topically. The term parenteral as used herein includes, subcutaneous, intravenous, intramuscular, intrasternal, infusion techniques or intraperitoneally.

The selection of the precise concentration, composition, and delivery regimen is influenced by, *inter alia*, the specific pharmacological properties of the particular selected compound, the intended use, the nature and severity of the condition being treated or diagnosed, the age, weight, gender, physical condition and mental acuity of the intended recipient as well as the route of administration. Such considerations are within the purview of the skilled artisan. Thus, the dosage regimen may vary widely, but can be determined routinely using standard methods.

Polyamides of the present invention are also useful for detecting the presence of double stranded DNA of a specific sequence for diagnostic or preparative purposes. The sample containing the double stranded DNA can be contacted by polyamide linked to a solid substrate, thereby isolating DNA comprising a desired sequence. Alternatively, polyamides linked to a suitable detectable marker, such as biotin, a hapten, a radioisotope or a dye molecule, can be contacted by a sample containing double stranded DNA.

The design of bifunctional sequence specific DNA binding molecules requires the integration of two separate entities: recognition and functional activity. Polyamides that specifically bind with subnanomolar affinity to the minor groove of a predetermined sequence of double stranded DNA are linked to a functional molecule, providing the corresponding bifunctional conjugates useful in molecular biology, genomic sequencing, and human medicine. Polyamides of this invention can be conjugated to a variety of functional molecules, which can be independently chosen from but is not limited to arylboronic acids, biotins, polyhistidines comprised from about 2 to 8 amino acids, haptens to which an antibody binds, solid phase supports, oligodeoxynucleotides, N-ethylnitrosourea, fluorescein, bromoacetamide, iodoacetamide, DL- $\alpha$ -lipoic acid, acridine, captothesin, pyrene, mitomycin, texas red, anthracene, anthrnilic acid, avidin, DAPI, isosulfan blue, malachite green, psoralen, ethyl red, 4-(psoraen-8-yloxy)-butyrate, tartaric acid, (+)- $\alpha$ -tocopheral, psoralen, EDTA, methidium, acridine, Ni(II)•Gly-Gly-His, thiazole orange (TO), Dansyl, pyrene, N-bromoacetamide, and gold particles. Such bifunctional polyamides are useful for DNA affinity capture, covalent DNA modification, oxidative DNA cleavage, DNA photocleavage. Such bifunctional polyamides are useful for DNA detection by providing a polyamide linked to a detectable label. Detailed instructions for synthesis of such bifunctional polyamides can be found in copending U.S. provisional application 60/043,444, the teachings of which are incorporated by reference.

DNA complexed to a labeled polyamide can then be determined using the appropriate detection system as is well known to one skilled in the art. For example, DNA associated with a polyamide linked to biotin can be detected by a streptavidin / alkaline phosphatase system.

The present invention also describes a diagnostic system, preferably in kit form, for assaying for the presence of the double stranded DNA sequence bound by the

polyamide of this invention in a body sample, such brain tissue, cell suspensions or tissue sections, or body fluid samples such as CSF, blood, plasma or serum, where it is desirable to detect the presence, and preferably the amount, of the double stranded DNA sequence bound by the polyamide in the sample according to the diagnostic methods described herein.

The diagnostic system includes, in an amount sufficient to perform at least one assay, a specific polyamide as a separately packaged reagent. Instructions for use of the packaged reagent(s) are also typically included. As used herein, the term "package" refers to a solid matrix or material such as glass, plastic (e.g., polyethylene, polypropylene or polycarbonate), paper, foil and the like capable of holding within fixed limits a polyamide of the present invention. Thus, for example, a package can be a glass vial used to contain milligram quantities of a contemplated polyamide or it can be a microliter plate well to which microgram quantities of a contemplated polyamide have been operatively affixed, i.e., linked so as to be capable of being bound by the target DNA sequence. "Instructions for use" typically include a tangible expression describing the reagent concentration or at least one assay method parameter such as the relative amounts of reagent and sample to be admixed, maintenance time periods for reagent or sample admixtures, temperature, buffer conditions and the like. A diagnostic system of the present invention preferably also includes a detectable label and a detecting or indicating means capable of signaling the binding of the contemplated polyamide of the present invention to the target DNA sequence. As noted above, numerous detectable labels, such as biotin, and detecting or indicating means, such as enzyme-linked (direct or indirect) streptavidin, are well known in the art.

Trauger, et al. (*Nature*, 382: 559-561) and Swalley, et al. (*J. Am. Chem. Soc.* 119: 6953-6961) have described recognition of DNA by certain polyamides at subnanomolar concentrations. Pairing specific carboxyamide groups allows for recognition of specific DNA sequences (Swalley, et al. *supra*). Polyamides comprising Hp, Im, and Py provide for coded targeting of pre-determined DNA sequences with high affinity and specificity. Hp, Im, and Py polyamides may be combined to form Im/Py, Py/Im, Hp/Py, and Py/Hp binding pairs which complement the four Watson-Crick base pairs A, C, G, and T. Table 1 illustrates such pairings.

**TABLE 1**

*Pairing Codes for Base Pair Recognition\**



Pair	G•C	C•G	T•A	A•T
Im/Py	+	-	-	-
Py/Im	-	+	-	-
Hp/Py	-	-	+	-
Py/Hp	-	-	-	+





\*favored (+), disfavored (-)

Three-, four-, five- or six-ring improved polyamides of the present invention are covalently coupled to form six-, eight-, ten- or twelve-ring structures, respectively, that bind specifically to four or six base pair targets, respectively, at subnanomolar concentrations. As such, the improved polyamides of the present invention may be directed to any DNA sequence comprised of A, C, G, or T.

The improved polyamides of the present invention comprise those having at least three consecutive carboxamide pairings for binding DNA in the minor groove of a regulatory sequence of a duplex gene sequence and a chiral hairpin turn with a stereochemical center substituted at the  $\gamma$ -position of the chiral hairpin turn of the molecule with the R-enantiomer of 2,4-diaminobutyric acid ( $H_2NHCHCH_2CHNH_2COOH$ ; "(R) $H_2N \gamma$ "). In addition, the present invention provides a methodology for covalently linking existing hairpin motifs without compromising DNA-binding and sequence specificity. The present invention provides improved polyamides for binding the minor groove of DNA to affect gene expression. Preferably, the bound polyamide inhibits gene expression.

The present invention comprises improved polyamides having three or four-ring polyamide structures covalently coupled to form six-, eight-, ten- or twelve-ring hairpin structures, respectively, of the general structures I-VIII:





where  $X_{1-12}$  and  $X'_{1-12}$  are independently an imidazole such as N-methylimidazole (Im), a pyrrole such as N-methylpyrrole (Py), or a hydroxypyrrole such as 3-hydroxy-N-methylpyrrole (Hp). In addition, an improved polyamide of the present invention may further include a aliphatic amino acid such as  $\beta$ -alanine residue ( $\beta$ ), an amide group such as dimethylaminopropylamide (Dp), an alcohol such as EtOH, an acid such as EDTA, or any derivative thereof may be joined to the  $\beta$  residue.

$\beta$ -alanine may also be utilized in place of a pyrrole amino acid in Formulas I-VIII. The use of  $\beta$ -alanine in place of a pyrrole or hydroxypyrrole amino acid in the synthetic methods provides aromatic/aliphatic pairing (Im/ $\beta$ ,  $\beta$ /Im, Py/ $\beta$ , and  $\beta$ /Py) and aliphatic/aliphatic pairing ( $\beta$ / $\beta$ ) substitution. Such substitutions may comprise those described in provisional application 60/042,022, incorporated herein by reference. The use of  $\gamma$ -aminobutyric acid, or a substituted  $\gamma$ -aminobutyric acid such as (R)-2,4-diaminobutyric acid, provides for preferred hairpin turns. Many other groups suitable for the purposes of practicing this invention are well known and widely available to one skilled in the art.

The polyamide subunit structures I-VIII above are covalently coupled through the  $\gamma$  residue which represents a  $-\text{NH}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CONH}-$  hairpin linkage derived from  $\gamma$ -aminobutyric acid or a chiral hairpin linkage derived from R-2,4-diaminobutyric acid. The present invention provides the reagents and methodologies for substituting the  $\gamma$ -residue of certain polyamides with a moiety such as (R)-2,4,-diaminobutyric acid ((R)<sup>H<sub>2</sub>N</sup>  $\gamma$ ). The NMR structure of a hairpin polyamide of sequence composition ImPyPy- $\gamma$ -PyPyPy complexed with a 5'-TGTTA-3' target site indicated that it was possible to substitute the  $\alpha$ -position of the  $\gamma$ -aminobutyric acid residue within the hairpin-DNA complex (de Claire, et al. *J. Am. Chem. Soc.* 1997, 119, 7909). Modeling indicated that replacing the  $\alpha$ -H of  $\gamma$  with an amino group that may confer an R-configuration at the  $\alpha$ -carbon could be accommodated within the floor and walls of the minor groove as demonstrated in Figure 1 and 2A. In contrast, the (S)-2,4,-diaminobutyric acid ((S)<sup>H<sub>2</sub>N</sup>  $\gamma$ ) linked hairpin is predicted to clash with the walls of the minor groove of the DNA helix as illustrated in Figures 1 and 2B.

In Formulas V-VIII, L represents an amino acid linking group such as  $\beta$ -alanine or 5-aminovaleric acid ( $\delta$ ) bound to the  $\gamma$  residue of a first polyamide and to the carboxytail of a second polyamide. As such, two or more polyamides may be linked, forming a tandemly-linked polyamide. Such a polyamide is said to be tandemly-linked or a tandem-linked polyamide.

P represents from zero to ten polyamides of formulas I-VIII that may be tandemly linked to the second polyamide. Preferably, P represents from zero to eight polyamids of formulas I-VIII. More preferably, P represents from zero to six polyamids of formulas I-VIII. More preferably, P represents from zero to four polyamides of formulas I-VIII. Most preferably, P represents from zero to two polyamides of formulas I-VIII. Tandem linking of polyamides provides expanded binding site size and increased binding affinity without compromising selectivity. Many other groups suitable for the purposes of practicing this invention are well known and widely available to one skilled in the art.

Baird, et al. (*J. Am. Chem. Soc.* 118: 6141-6146) and PCT/US97/003332 describe methods for synthesis of polyamides which are suitable for preparing polyamides of this invention. Polyamides of the present invention may be synthesized by solid phase

methods using compounds such as Boc-protected 3-methoxypyrrole, imidazole, and pyrrole aromatic amino acids, which are cleaved from the support by aminolysis, deprotected with sodium thiophenoxide, and purified by reverse-phase HPLC. The identity and purity of the polyamides may be verified using any of a variety of analytical techniques available to one skilled in the art such as  $^1\text{H-NMR}$ , analytical HPLC, and/or matrix-assisted laser-desorption ionization time-of-flight mass spectrometry (MALDI-TOF MS-monoisotopic).

Described herein is the synthesis of a new class of chiral hairpin polyamides and their characterization with regard to DNA binding affinity and sequence specificity. The present invention provides one skilled in the art with the reagents and methodologies for substitution of the prochiral  $\gamma$ -turn with either enantiomer of 2,4-diaminobutyric acid. In addition, the invention provides the dicationic six-ring enantiomeric polyamides (+)-ImPyPy-(R) $^{\text{H}_2\text{N}}$   $\gamma$ -PyPyPy- $\beta$ -Dp (1-R) and (-)-ImPyPy-(S) $^{\text{H}_2\text{N}}$   $\gamma$ -PyPyPy- $\beta$ -Dp (1-S) which may be synthesized by solid phase methods. In certain experiments, the monocationic polyamide (+)-ImPyPy-(R) $^{\text{H}_2\text{N}}$   $\gamma$  PyPyPy- $\beta$ -EtOH (2-R), which lacks a charge at the C-terminus, may be prepared and utilized as a control. To further study steric effects, the  $\gamma$ -acetamido polyamides (+)-ImPyPy-(R) $^{\text{Ac}}$   $\gamma$ -PyPyPy- $\beta$ -Dp (3-R) and (-)-ImPyPy-(S) $^{\text{Ac}}$   $\gamma$ -PyPyPy- $\beta$ -Dp (3-S) may be utilized (Figure 3; Baird, et al. 1996. *J. Am. Chem. Soc.* 118: 6141). The present invention further provides the EDTA analogs ImPyPy-(R) $^{\text{H}_2\text{N}}$   $\gamma$ -PyPyPy- $\beta$ -Dp-EDTA $\cdot\text{Fe(II)}$  (4-R $\cdot\text{Fe(II)}$ ), ImPyPy-(S) $^{\text{H}_2\text{N}}$   $\gamma$ -PyPyPy- $\beta$ -Dp-EDTA $\cdot\text{Fe(II)}$  (4-S $\cdot\text{Fe(II)}$ ); ImPyPy-(R) $^{\text{EDTA}\cdot\text{Fe(II)}}$   $\gamma$ -PyPyPy- $\beta$ -Dp (5-R $\cdot\text{Fe(II)}$ ), and ImPyPy-(S) $^{\text{EDTA}\cdot\text{Fe(II)}}$   $\gamma$ -PyPyPy- $\beta$ -Dp (5-S $\cdot\text{Fe(II)}$ ) that may be utilized to confirm the binding orientation of the modified hairpins at specific DNA binding sites (Figure 3).

Tandemly-linked polyamides of the present invention are also provided by the instant invention. The primary turn-amino group provides a potential site for covalently tethering two hairpins. In one potential linkage arrangement, the C-terminus of the first hairpin is coupled to the  $\alpha$ -amino group of the  $\gamma$ -turn of the second amino acid linker. The present invention provides twelve ting polyamides exemplified by ImPyPy-(R)[ImPyPy-(R) $^{\text{H}_2\text{N}}$   $\gamma$ -PyPyPy- $\beta$ ] $^{\text{HN}}$   $\gamma$ -PyPyPy- $\beta$ -Dp and ImPyPy-(R)[ImPyPy-(R) $^{\text{H}_2\text{N}}$   $\gamma$ -PyPyPy- $\beta$ ] $^{\text{HN}}$   $\gamma$ -PyPyPy- $\beta$ -Dp (Figure 14). The DNA binding properties of certain

polyamides of the present invention were determined on a series of DNA fragments containing 10, 11 and 12 base pair target sites. The present invention further provides an exemplary affinity cleaving derivative  $\text{ImPyPy-(R)[ImPyPy-(R)}^{\text{EDTA}}\gamma\text{PyPyPy-}\delta]^{\text{HN}}\gamma\text{PyPyPy-}\beta\text{-Dp}$ , which was utilized to confirm a single predicted binding orientation for the tandemly-linked polyamide. Methodologies for the determination of the DNA-binding affinity and sequence selectivity of tandem improved polyamides is also provided.

The present invention reveals to one skilled in the art properties of chiral structure elements that may be utilized as a guide in the design of more efficient polyamides. For instance, the present invention provides amine substituents on the  $(R)^{\text{H}}_2\text{N}\gamma$  turn amino acid that enhance the DNA binding affinity and specificity relative to the unsubstituted parent hairpin, providing for an optimized class of hairpin polyamides. Also provided are acetamido substituents at the  $(R)^{\text{H}}_2\text{N}\gamma$  that do not compromise affinity or specificity relative to the parent hairpin, providing for a convenient synthetic attachment point at the 'capped' end of the molecule. In addition, the invention described herein provides  $(S)^{\text{H}}_2\text{N}\gamma$ -linked hairpins that bind with enhanced affinity to reverse orientation sites relative to the parent hairpin and  $(R)^{\text{H}}_2\text{N}\gamma$ -linked hairpins that bind with enhanced specificity relative to the parent hairpin indicating that  $\gamma$ -turn substituents may regulate hairpin polyamide binding orientational preference. The invention further provides the skilled artisan with the necessary tools and methodologies for developing tandemly-linked polyamides to increase the polyamide binding site size, and increase affinity without compromising sequence selectivity.

The examples listed above and those illustrated below represent only certain embodiments of the present invention and are not limiting of the specification and claims in any way.



## EXAMPLES

### Example 1

#### Synthesis of Improved Polyamides

Two polyamide-resins ImPyPy-(R)<sup>Fmoc</sup>γ-PyPyPy-β-Pam-resin and ImPyPy-(S)<sup>Fmoc</sup>γ-PyPyPy-β-Pam-resin, were synthesized in 14 steps from Boc-β-alanine-Pam-resin (1 g resin, 0.2 mmol/g substitution) using previously described Boc-chemistry machine-assisted protocols (Figure 4; Baird, et al. *J. Am. Chem. Soc.* **1996**, *118*, 6141). (R)- and (S)-2,4-diaminobutyric acid residues were introduced as orthogonally protected N-γ-Fmoc-N-γ-Boc derivatives (HBTU, DIEA). Fmoc protected polyamide resins ImPyPy-(R)<sup>Fmoc</sup>γ-PyPyPy-β-Pam-resin and ImPyPy-(S)<sup>Fmoc</sup>γ-PyPyPy-β-Pam-resin were treated with 1:4 DMF:Piperidine (22 °C, 30 min.) to provide ImPyPy-(R)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Pam-resin and ImPyPy-(S)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Pam-resin, respectively. A single-step aminolysis of the resin ester linkage was used to cleave the polyamide from the solid support. A sample of resin (240 mg) was treated with either dimethylaminopropylamine (55 °C, 18 h) to provide 1-R, 1-S, 3-R, and 3-S or ethanolamine (55 °C, 18 h) to provide 2-R. Resin cleavage products were purified by reverse phase HPLC to provide ImPyPy-(R)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp (1-R), ImPyPy-(S)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp (1-S), and ImPyPy-(R)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-EtOH (2-R). The stereochemical purity of 1-R was determined to be > 98% by Mosher amide analysis (Dale, et al. *J. Am. Chem. Soc.* **1973**, *95*, 512; Yamaguchi, et al. *Asymmetric Synthesis (Vol. 1), Analytical Methods* p.125-152, J. D. Morrison (ed.) Academic Press (1983)). 1-R,R and 1-R,S Mosher amides were prepared by reaction of 1-R with HOBt activated esters generated *in situ* from (R)-α-methoxy-α-(trifluoromethyl)phenylacetic acid and (S)-α-methoxy-α-(trifluoromethyl)phenylacetic acid. For synthesis of analogs modified with EDTA at the carboxy-terminus, the amine-resin was treated with Boc-anhydride (DMF, DIEA, 55 °C, 30 min) to provide ImPyPy-(R)<sup>Boc</sup>γ-PyPyPy-β-Pam-resin and ImPyPy-(S)<sup>Boc</sup>γ-PyPyPy-β-Pam-resin (Figure 4). A sample of Boc-resin was then cleaved with 3,3'-diamino-N-methyldipropylamine (55 °C,

18 h) and purified by reversed phase HPLC to provide either ImPyPy-(R)<sup>Boc</sup>γ-PyPyPy-β-Dp-NH<sub>2</sub> (**1-R-Boc-NH<sub>2</sub>**) or ImPyPy-(S)<sup>Boc</sup>γ-PyPyPy-β-Dp-NH<sub>2</sub> (**1-S-Boc-NH<sub>2</sub>**) which afford free primary amine groups at the C-terminus suitable for post-synthetic modification. The polyamide-amines **1-R-Boc-NH<sub>2</sub>** and **1-S-Boc-NH<sub>2</sub>** were treated with an excess of EDTA-dianhydride (DMSO/NMP, DIEA, 55 °C, 15 min) and the remaining anhydride hydrolyzed (0.1 M NaOH, 55 °C, 10 min). The Boc protected EDTA modified polyamides ImPyPy-(R)<sup>Boc</sup>γ-PyPyPy-β-Dp-EDTA (**4-R-Boc**) and ImPyPy-(S)<sup>Boc</sup>γ-PyPyPy-β-Dp-EDTA (**4-S-Boc**) were isolated by HPLC. Individual Boc-EDTA-polyamides were deprotected with neat TFA (22 °C, 1 h) to provide the respective C-terminal EDTA derivatives, ImPyPy-(R)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp-EDTA (**4-R**) and ImPyPy-(S)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp-EDTA (**4-S**). For the synthesis of acetamide-turn or EDTA-turn derivatives, a sample of the γ-amino polyamide ImPyPy-(R)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp (**1-R**) or ImPyPy-(S)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp (**1-S**) was treated with an excess of either acetic anhydride or EDTA-dianhydride (DMSO/NMP, DIEA 55 °C, 30 min) and the remaining anhydride hydrolyzed (0.1 M NaOH, 55 °C, 10 min). The polyamides ImPyPy-(R)<sup>Ac</sup>γ-PyPyPy-β-Dp (**3-R**), ImPyPy-(S)<sup>Ac</sup>γ-PyPyPy-β-Dp (**3-S**), ImPyPy-(R)<sup>EDTA</sup>γ-PyPyPy-β-Dp (**5-R**) and ImPyPy-(S)<sup>EDTA</sup>γ-PyPyPy-β-Dp (**5-S**) were then isolated by reverse phase HPLC. The six-ring hairpin polyamides described here are soluble in aqueous solution at concentrations 10 mM at 37°C.

#### A. Materials

Dicyclohexylcarbodiimide (DCC), Hydroxybenzotriazole (HOBt), 2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexa-fluorophosphate (HBTU) and 0.2 mmol/gram Boc-β-alanine-(-4-carboxamidomethyl)-benzyl-ester-copoly(styrene-divinylbenzene) resin (Boc-γ-Pam-Resin) was purchased from Peptides International (0.2 mmol/gram) (R)-2-Fmoc-4-Boc-diaminobutyric acid, (S)-2-Fmoc-4-Boc-diaminobutyric acid, and (R)-2-amino-4-Boc-diaminobutyric acid were from Bachem. N,N-diisopropylethylamine (DIEA), N,N-dimethylformamide (DMF), N-methylpyrrolidone

(NMP), DMSO/NMP, Acetic anhydride ( $\text{Ac}_2\text{O}$ ), and 0.0002 M potassium cyanide/pyridine were purchased from Applied Biosystems. Dichloromethane (DCM) and triethylamine (TEA) were reagent grade from EM, thiophenol (PhSH), dimethylaminopropylamine (Dp), (*R*)- $\alpha$ -methoxy- $\alpha$ -(trifluoromethyl)phenylacetic acid ((*R*)MPTA) and (*S*)- $\alpha$ -methoxy- $\alpha$ -(trifluoromethyl)phenylacetic acid ((*S*)MPTA) were from Aldrich, trifluoroacetic acid (TFA) Biograde from Halocarbon, phenol from Fisher, and ninhydrin from Pierce. All reagents were used without further purification.

Quik-Sep polypropylene disposable filters were purchased from Isolab Inc. A shaker for manual solid phase synthesis was obtained from St. John Associates, Inc. Screw-cap glass peptide synthesis reaction vessels (5 mL and 20 mL) with a #2 sintered glass frit were made as described by Kent (*Annu. Rev. Biochem.* 1988, 57, 957).  $^1\text{H}$  NMR spectra were recorded on a General Electric-QE NMR spectrometer at 300 MHz with chemical shifts reported in parts per million relative to residual solvent. UV spectra were measured in water on a Hewlett-Packard Model 8452A diode array spectrophotometer. Optical rotations were recorded on a JASCO Dip 1000 Digital Polarimeter. Matrix-assisted, laser desorption/ionization time of flight mass spectrometry (MALDI-TOF) was performed at the Protein and Peptide Microanalytical Facility at the California Institute of Technology. HPLC analysis was performed on either a HP 1090M analytical HPLC or a Beckman Gold system using a RAINEN  $\text{C}_{18}$ , Microsorb MV,  $5\mu\text{m}$ ,  $300 \times 4.6$  mm reversed phase column in 0.1% (wt/v) TFA with acetonitrile as eluent and a flow rate of 1.0 mL/min, gradient elution 1.25% acetonitrile/min. Preparatory reverse phase HPLC was performed on a Beckman HPLC with a Waters DeltaPak  $25 \times 100$  mm,  $100\mu\text{m}$   $\text{C}_{18}$  column equipped with a guard, 0.1% (wt/v) TFA, 0.25% acetonitrile/min. Distilled water was obtained from a Millipore MilliQ water purification system, and all buffers were  $0.2\mu\text{m}$  filtered.

Enzymes were purchased from Boehringer-Mannheim and used with their supplied buffers. Deoxyadenosine and thymidine  $5'-[\gamma^{32}\text{P}]$  triphosphates were obtained

from Amersham, and deoxyadenosine 5'-[ $\gamma$ - $^{32}$ P]triphosphate was purchased from I.C.N. Sonicated, deproteinized calf thymus DNA was acquired from Pharmacia. RNase free water was obtained from USB and used for all footprinting reactions. All other reagents and materials were used as received. All DNA manipulations were performed according to standard protocols (Sambrook, J.; Fritsch, E.F.; Maniatis, T. *Molecular Cloning*; Cold Spring Harbor Laboratory: Cold Spring Harbor, NY, 1989).

### B. ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (1-R)

ImPyPy-(R)<sup>Fmoc</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-Resin was synthesized in a stepwise fashion by machine-assisted solid phase methods (Baird, et al. *J. Am. Chem. Soc.* **1996**, *118*, 6141). (R)-2-Fmoc-4-Boc-diaminobutyric acid (0.7 mmol) was incorporated as previously described for Boc- $\gamma$ -aminobutyric acid. ImPyPy-(R)<sup>Fmoc</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-Resin was placed in a glass 20 mL peptide synthesis vessel and treated with DMF (2 mL), followed by piperidine (8 mL) and agitated (22 °C, 30 min.). ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-resin was isolated by filtration, and washed sequentially with an excess of DMF, DCM, MeOH, and ethyl ether and the amine-resin dried *in vacuo*. A sample of ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-resin (240 mg, 0.18 mmol/gram) was treated with neat dimethylaminopropylamine (2 mL) and heated (55 °C) with periodic agitation for 16 h. Resin substitution can be calculated as  $L_{\text{new}}(\text{mmol/g}) = L_{\text{old}} / (1 + L_{\text{old}}(W_{\text{new}} - W_{\text{old}}) \times 10^{-3})$ , where L is the loading (mmol of amine per gram of resin), and W is the weight (gmol<sup>-1</sup>) of the growing polyamide attached to the resin (Barlos, et al. *Int. J. Peptide Protein Res.* **1991**, *37*, 513). The reaction mixture was placed in an oven and periodically agitated (55 °C, 16 h). The reaction mixture was then filtered to remove resin, 0.1% (wt/v) TFA added (6 mL) and the resulting solution purified by reversed phase HPLC. ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ -Dp is recovered upon lyophilization of the appropriate fractions as a white powder (32 mg, 66% recovery).  $[\alpha]_{\text{D}}^{20} +14.6$  (c 0.05, H<sub>2</sub>O); UV (H<sub>2</sub>O) $\lambda_{\text{max}}$  246, 310 (50,000); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) 10.56 (s, 1 H), 10.47 (s, 1 H), 9.97 (s,

1 H), 9.94 (s, 1 H), 9.88 (s, 1 H), 9.4 (br s, 1 H), 8.28 (s, 3 H), 8.22 (m, 1 H), 8.03 (m, 2 H), 7.38 (s, 1 H), 7.25 (d, 1 H,  $J = 1.6$  Hz), 7.22 (d, 1 H,  $J = 1.5$  Hz), 7.19 (d, 1 H,  $J = 1.5$  Hz), 7.16 (d, 1 H,  $J = 1.6$  Hz), 7.14 (d, 1 H,  $J = 1.8$  Hz), 7.12 (d, 1 H,  $J = 1.7$  Hz), 7.03 (m, 2 H), 6.95 (d, 1 H,  $J = 1.6$  Hz), 6.91 (d, 1 H,  $J = 1.6$  Hz), 6.85 (d, 1 H,  $J = 1.6$  Hz), 3.96 (s, 3 H), 3.83 (s, 3 H), 3.81 (m, 6 H), 3.79 (s, 3 H), 3.76 (s, 3 H), 3.33 (q, 2 H,  $J = 6.3$  Hz), 3.25 (q, 2 H,  $J = 5.7$  Hz), 3.05 (q, 2 H,  $J = 5.9$  Hz), 2.96 (q, 2 H,  $J = 5.3$  Hz), 2.71 (d, 6 H,  $J = 4.9$  Hz), 2.32 (t, 2 H,  $J = 7.1$  Hz), 1.95 (q, 2 H,  $J = 5.9$  Hz), 1.70 (quintet, 2 H,  $J = 7.3$  Hz); MALDI-TOF-MS (monoisotopic), 992.5 (992.5 calc. for  $C_{47}H_{62}N_{17}O_8$ ).

A hydrogen bonding model of the 1:1 polyamide:DNA complex formed between the hairpin polyamide ImPyPy-(*R*)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp (**1-R**) with a 5'-TGTTA-3' site is illustrated in Figure 1A. Lone pairs of N<sub>3</sub> of purines and O<sub>2</sub> of pyrimidines are shown as circles with dots. The N<sub>2</sub> hydrogen of guanine is indicated by circles containing an "H" and putative hydrogen bonds are illustrated by dotted lines. A schematic binding model is also demonstrated where the imidazole and pyrrole rings are represented as shaded and unshaded spheres, respectively, and the β-alanine residue is represented as an unshaded diamond. Figure 2 shows models derived from the NMR structure coordinates of ImPyPy-γ-PyPyPy-β-Dp•5'-TGTTA-3' using InsightII software. Figure 3 shows the structure of the 2-*R* polyamides.

### C. ImPyPy-(*R*)<sup>(R)MTPA</sup>γ-PyPyPy-β-Dp (**1-R,R**)

(*R*)-α-methoxy-α-(trifluoromethyl)phenylacetic acid (117 mg, 0.5 mmol) and HOBt (70 mg, 0.5 mmol) were dissolved in DMF (1 mL), DCC (100 mg, 0.5 mmol) added and the solution agitated for 30 min at 22 °C. A sample of the activated ester solution (100 μL, 0.05 mmol) was added to ImPyPy-(*R*)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp **1-R** (10 mg, 0.01 mmol), DIEA (50 μL) added, and the solution agitated for 3h (22 °C). DMF (1 mL) followed by 0.1% (wt/v) TFA (6 mL) was then added to the reaction mixture and the resulting solution purified by reversed phase HPLC (1% acetonitrile/min.) under

conditions which were determined to separate the diastereomers. ImPyPy-(R)<sup>(R)MTPA</sup>γ-PyPyPy-β-Dp is recovered as a white powder upon lyophilization of the appropriate fractions (6 mg, 53% recovery). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ10.50 (s, 1 H), 10.14 (s, 1 H), 9.92 (s, 2 H), 9.88 (s, 1 H), 9.2 (br s, 1 H), 8.43 (d, 1 H, *J* = 7.0 Hz), 8.02 (m, 3 H), 7.92 (m, 1 H), 7.47 (m, 2 H), 7.41 (m, 2 H), 7.36 (s, 1 H), 7.24 (m, 1 H), 7.19 (m, 1 H), 7.15 (m, 1 H), 7.12 (m, 3 H), 7.01 (m, 2 H), 6.90 (m, 3 H), 6.83 (m, 1 H), 4.46 (q, 1 H, *J* = 5.5 Hz), 3.94 (s, 3 H), 3.79 (m, 9 H), 3.75 (m, 6 H), 3.32 (m, 4 H), 3.05 (m, 2 H), 2.94 (m, 2 H), 2.68 (d, 6H, *J* = 4.0 Hz), 2.28 (t, 2 H, *J* = 6.3 Hz), 1.93 (q, 2 H, *J* = 6.1 Hz), 1.66 (quintet, 2 H, *J* = 6.0 Hz), 1.18 (s, 3 H); MALDI-TOF-MS (monoisotopic), 1208.5 (1208.5 calc. for C<sub>57</sub>H<sub>68</sub>F<sub>3</sub>N<sub>17</sub>O<sub>10</sub>).

#### D. ImPyPy-(R)<sup>(S)MTPA</sup>γ-PyPyPy-β-Dp 1-*R,S*

ImPyPy-(R)<sup>(S)MTPA</sup>γ-PyPyPy-β-Dp was prepared from (*S*)-α-methoxy-α-(trifluoromethyl)phenylacetic acid as described for 1-*R,R* (5 mg, 45% recovery). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ10.47 (s, 1 H), 10.08 (s, 1 H), 9.92 (s, 2 H), 9.88 (s, 1 H), 9.2 (br s, 1 H), 8.43 (d, 1 H, *J* = 6.9 Hz), 8.02 (m, 3 H), 7.46 (m, 2 H), 7.40 (m, 2 H), 7.36 (s, 1 H), 7.23 (m, 1 H), 7.19 (m, 1 H), 7.14 (m, 1 H), 7.12 (m, 3 H), 7.01 (m, 2 H), 6.87 (m, 3 H), 6.83 (m, 1 H), 4.44 (q, 1 H, *J* = 6.5 Hz), 3.94 (s, 3 H), 3.79 (m, 9 H), 3.75 (m, 6 H), 3.28 (m, 4 H), 3.06 (m, 4 H), 2.94 (m, 2 H), 2.69 (d, 6H, *J* = 4.5 Hz), 2.28 (t, 2 H, *J* = 6.5 Hz), 1.93 (q, 2 H, *J* = 6.1 Hz), 1.66 (quintet, 2 H, *J* = 6.0 Hz), 1.18 (s, 3 H); MALDI-TOF-MS (monoisotopic), 1209.0 (1208.5 calc. for C<sub>57</sub>H<sub>68</sub>F<sub>3</sub>N<sub>17</sub>O<sub>10</sub>).

#### E. ImPyPy-(S)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp (1-*S*)

ImPyPy-(S)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp was prepared as described for 1-*R* (23 mg, 49% recovery). [α]<sub>D</sub><sup>20</sup> -14.2 (c 0.04, H<sub>2</sub>O); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) identical to 1-*R*; MALDI-TOF-MS (monoisotopic), 992.5 (992.5 calc. for C<sub>47</sub>H<sub>62</sub>N<sub>17</sub>O<sub>8</sub>). Figure 2B illustrates a binding model for the ImPyPy-(S)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp (1-*S*) polyamide to the DNA sequence 5'-TGTTA-3'.

#### F. ImPyPy-(R)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-EtOH (2-*R*)

A sample of ImPyPy-(R)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Pam-resin (240 mg, 0.18 mmol/gram) was treated with neat ethanolamine (2 mL) and heated (55 °C) with periodic agitation for 16 h. The reaction mixture was then filtered to remove resin, 0.1% (wt/v) TFA added (6 mL) and the resulting solution purified by reversed phase HPLC to provide ImPyPy-(R)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-EtOH as a white powder upon lyophilization of the appropriate fractions (21 mg, 46% recovery).  $[\alpha]_D^{20} +18.6$  (c 0.04, H<sub>2</sub>O); UV (H<sub>2</sub>O)  $\lambda_{\max}$  246, 310 (50,000); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ10.55 (s, 1 H), 10.48 (s, 1 H), 9.97 (s, 1 H), 9.94 (s, 1 H), 9.89 (s, 1 H), 8.24 (m, 4 H), 8.00 (t, 1 H, *J* = 4.1 Hz), 7.89 (t, 1 H, *J* = 5.8 Hz), 7.38 (s, 1 H), 7.25 (d, 1 H, *J* = 1.6 Hz), 7.22 (d, 1 H, *J* = 1.6 Hz), 7.21 (d, 1 H, *J* = 1.5 Hz), 7.16 (m, 2 H), 7.14 (d, 1 H, *J* = 1.6 Hz), 7.03 (d, 1 H, *J* = 1.7 Hz), 6.99 (d, 1 H, *J* = 1.4 Hz), 6.95 (d, 1 H, *J* = 1.6 Hz), 6.91 (d, 1 H, *J* = 1.5 Hz), 6.78 (d, 1 H, *J* = 1.5 Hz), 5.33 (m, 1 H), 3.95 (s, 3 H), 3.83 (s, 3 H), 3.81 (m, 6 H), 3.79 (s, 3 H), 3.76 (s, 3 H), 3.37 (q, 2 H, *J* = 6.2 Hz), 3.07 (q, 2 H, *J* = 5.9 Hz), 2.29 (t, 2 H, *J* = 7.1 Hz), 1.93 (q, 2 H, *J* = 5.8 Hz), 1.20 (m, 4 H); MALDI-TOF-MS (monoisotopic), 951.4 (951.4 calc. for C<sub>44</sub>H<sub>55</sub>N<sub>16</sub>O<sub>9</sub>). The 2-*R* polyamide is shown in Figure 3.

#### G. ImPyPy-(R)<sup>Ac</sup>γ-PyPyPy-β-Dp (3-*R*)

A sample of ImPyPy-(R)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp (4 mg) in DMSO (1 mL) was treated with a solution of acetic anhydride (1 mL) and DIEA (1 mL) in DMF (1 mL) and heated (55 °C) with periodic agitation for 30 min. Residual acetic anhydride was hydrolyzed (0.1 M NaOH, 1 mL, 55 °C, 10 min.), 0.1% (wt/v) TFA was added (6 mL) and the resulting solution purified by reversed phase HPLC to provide ImPyPy-(R)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp is recovered as a white powder upon lyophilization of the appropriate fractions (2 mg, 50% recovery).  $[\alpha]_D^{20} +20.5$  (c 0.06, H<sub>2</sub>O); UV (H<sub>2</sub>O)  $\lambda_{\max}$  242, 304 (50,000); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ10.49 (s, 1 H), 10.06 (s, 1 H), 9.94 (m, 2 H), 9.00 (s, 1 H), 9.4 (br s, 1 H), 8.21 (d, 1 H, *J* = 7.8 Hz), 8.06 (m, 2 H), 8.00 (t, 1 H, *J* = 6.2 Hz), 7.39 (s, 1 H), 7.27 (d, 1 H, *J* = 1.7 Hz), 7.21 (d, 1 H, *J* = 1.6 Hz), 7.18 (m, 2 H), 7.14 (m, 2 H), 7.03 (m, 2 H), 6.90 (d, 1 H, *J* = 1.6 Hz), 6.86 (m, 2 H), 4.43 (q, 1 H, *J* = 7.5 Hz), 3.96 (s, 3 H), 3.82

(m, 9 H), 3.73 (m, 6 H), 3.37 (q, 2 H,  $J = 5.8$  Hz), 3.11 (q, 2 H,  $J = 6.9$  Hz), 2.98 (q, 2 H,  $J = 5.4$  Hz), 2.79 (q, 2 H,  $J = 5.3$  Hz), 2.71 (d, 6 H,  $J = 4.7$  Hz), 2.33 (t, 2 H,  $J = 6.2$  Hz), 1.97 (s, 3 H), 1.70 (quintet, 2 H,  $J = 6.0$  Hz) MALDI-TOF-MS (average), 1035.1 (1035.2 calc. for M+H). The 3-*R* polyamide is shown in Figure 3.

5

#### H. ImPyPy-(S)<sup>Ac</sup>γPyPyPy-β-Dp (3-*S*)

ImPyPy-(S)<sup>Ac</sup>γPyPyPy-β-Dp was prepared as described for 3-*R*. (2 mg, 50% recovery).  $[\alpha]_D^{20} -16.4$  (c 0.07, H<sub>2</sub>O); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) is identical to 3-*R*; MALDI-TOF-MS (monoisotopic), 1034.6 (1034.5 calc. for C<sub>49</sub>H<sub>64</sub>N<sub>17</sub>O<sub>9</sub>).

#### I. ImPyPy-(R)<sup>Boc</sup>γPyPyPy-β-Dp-NH<sub>2</sub> (4-*R*-Boc-NH<sub>2</sub>)

A sample of ImPyPy-(R)<sup>H<sub>2</sub>N</sup>γPyPyPy-β-Pam-resin (300 mg, 0.18 mmol/gram). Resin substitution can be calculated as  $L_{\text{new}}(\text{mmol/g}) = L_{\text{old}} / (1 + L_{\text{old}}(W_{\text{new}} - W_{\text{old}}) \times 10^{-3})$ , where *L* is the loading (mmol of amine per gram of resin), and *W* is the weight (gmol<sup>-1</sup>) of the growing polyamide attached to the resin. see: Barlos, et al. *Int. J. Peptide Protein Res.* 1991, 37, 513.) was treated a solution of Boc-anhydride (500 mg) and DIEA (1 mL) in DMF (4 ml) and heated (55 °C) with periodic agitation for 30 min. ImPyPy-(R)<sup>Boc</sup>γ-PyPyPy-β-Pam-resin was isolated by filtration, and washed sequentially with an excess of DMF, DCM, MeOH, and ethyl ether and the dried *in vacuo*. A sample of ImPyPy-(R)<sup>Boc</sup>γ-PyPyPy-β-Pam-resin (240 mg, 0.18 mmol/gram) was treated with neat 3,3'-diamino-*N*-methyldipropylamine (2 mL) and heated (55 °C) with periodic agitation for 16h. The reaction mixture was then filtered to remove resin, 0.1% (wt/v) TFA added (6 mL) and the resulting solution purified by reversed phase HPLC to provide ImPyPy-(R)<sup>Boc</sup>γ-PyPyPy-β-Dp-NH<sub>2</sub> as a white powder upon lyophilization of the appropriate fractions (18 mg, 36% recovery);  $[\alpha]_D^{20} -30$  (c 0.05, H<sub>2</sub>O); UV (H<sub>2</sub>O)  $\lambda_{\text{max}}$  240, 306 (50,000); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>)  $\delta$ 10.59 (s, 1 H), 10.16 (s, 1 H), 10.04 (m, 2 H), 10.00 (s,



1 H), 9.4 (br s, 1 H), 8.31 (d, 1 H,  $J = 7.8$  Hz), 8.16 (m, 2 H), 8.10 (t, 1 H,  $J = 6.2$  Hz), 7.89 (t, 1 H,  $J = 5.8$  Hz), 7.49 (s, 1 H), 7.37 (d, 1 H,  $J = 1.7$  Hz), 7.22 (d, 1 H,  $J = 1.6$  Hz), 7.21 (d, 1 H,  $J = 1.5$  Hz), 7.16 (m, 2 H), 7.14 (d, 1 H,  $J = 1.6$  Hz), 7.03 (d, 1 H,  $J = 1.7$  Hz), 6.99 (d, 1 H,  $J = 1.4$  Hz), 6.95 (d, 1 H,  $J = 1.6$  Hz), 6.91 (d, 1 H,  $J = 1.5$  Hz), 6.78 (d, 1 H,  $J = 1.5$  Hz), 5.33 (m, 1 H), 3.95 (s, 3 H), 3.83 (s, 3 H), 3.81 (m, 6 H), 3.79 (s, 3 H), 3.76 (s, 3 H), 3.37 (q, 2 H,  $J = 6.2$  Hz), 3.07 (q, 2 H,  $J = 5.9$  Hz), 2.29 (t, 2 H,  $J = 7.1$  Hz), 1.93 (q, 2 H,  $J = 5.8$  Hz), 1.20 (m, 4 H); MALDI-TOF-MS (monoisotopic), 1135.3 (1135.6 calc. for  $C_{54}H_{75}N_{18}O_{10}$ ).

**J. ImPyPy-(S)<sup>Boc</sup>- $\gamma$ -PyPyPy- $\beta$ -Dp-NH<sub>2</sub> (4-S-Boc-NH<sub>2</sub>)**

ImPyPy-(S)<sup>Boc</sup>- $\gamma$ -PyPyPy- $\beta$ -Dp-NH<sub>2</sub> was prepared as described for 4-R. (16 mg, 32% recovery).  $[\alpha]_D^{20}$  -30 (c 0.05, H<sub>2</sub>O); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) is identical to 4-R-Boc-NH<sub>2</sub>; MALDI-TOF-MS (monoisotopic), 1135.4 (1135.6 calc. for  $C_{54}H_{75}N_{18}O_{10}$ ).

**K. ImPyPy-(R)<sup>Bo</sup>- $\gamma$ -PyPyPy- $\beta$ -Dp-EDTA (4-R-Boc)**

Excess EDTA-dianhydride (50 mg) was dissolved in DMSO/NMP (1 mL) and DIEA (1 mL) by heating at 55 °C for 5 min. The dianhydride solution was added to ImPyPy-(S)<sup>Boc</sup>- $\gamma$ -PyPyPy- $\beta$ -Dp-NH<sub>2</sub> (10.4 mg, 10  $\mu$ mol) dissolved in DMSO (750  $\mu$ L). The mixture was heated (55 °C, 25 min.) and the remaining EDTA-anhydride hydrolyzed (0.1M NaOH, 3 mL, 55 °C, 10 min). Aqueous TFA (0.1% wt/v) was added to adjust the total volume to 8 mL and the solution purified directly by reversed phase HPLC to provide ImPyPy-(R)<sup>Boc</sup>- $\gamma$ -PyPyPy- $\beta$ -Dp-EDTA (4-R-Boc) as a white powder upon lyophilization of the appropriate fractions (4 mg, 40% recovery). MALDI-TOF-MS (monoisotopic), 1409.6 (1409.7 calc. for  $C_{64}H_{89}N_{20}O_{17}$ ).

**L. ImPyPy-(S)<sup>Boc</sup>- $\gamma$ -PyPyPy- $\beta$ -Dp-EDTA (4-S-Boc)**

ImPyPy-(S)<sup>Boc</sup>- $\gamma$ -PyPyPy- $\beta$ -Dp-NH<sub>2</sub> (12.0 mg, 12  $\mu$ mol) was converted to 4-S-Boc as described for 4-R-Boc (4 mg, 33% recovery). MALDI-TOF-MS (monoisotopic), 1409.7 (1409.7 calc. for  $C_{64}H_{89}N_{20}O_{17}$ ).

**M. ImPyPy-(R)<sup>H<sub>2</sub>N</sup>- $\gamma$ -PyPyPy- $\beta$ -Dp-EDTA (4-R)**

A sample of ImPyPy-(R)<sup>Boc</sup>γ-PyPyPy-β-Dp-EDTA (2.1 mg) in DMSO (750 μL) was placed in a 50 mL flask and treated with TFA (15 mL, 22 °C, 2 h). Excess TFA was removed *in vacuo*, water added (6 mL) and the resulting solution purified by reversed phase HPLC to provide ImPyPy-(R)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp-EDTA as a white powder upon lyophilization of the appropriate fractions (1.3 mg, 50% recovery). MALDI-TOF-MS (monoisotopic), 1309.5 (1309.6 calc. for C<sub>59</sub>H<sub>81</sub>N<sub>20</sub>O<sub>15</sub>).

**N. ImPyPy-(S)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp-EDTA (4-S)**

ImPyPy-(S)<sup>Boc</sup>γ-PyPyPy-β-Dp-EDTA (3.0 mg) was converted to 4-S as described for 4-R (1 mg, 33% recovery). MALDI-TOF-MS (monoisotopic), 1309.5 (1309.6 calc. for C<sub>59</sub>H<sub>81</sub>N<sub>20</sub>O<sub>15</sub>).

**O. ImPyPy-(R)<sup>EDTA</sup>γ-PyPyPy-β-Dp (5-R)**

Excess EDTA-dianhydride (50 mg) was dissolved in DMSO/NMP (1 mL) and DIEA (1 mL) by heating at 55 °C for 5 min. The dianhydride solution was added to ImPyPy-(R)<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp (1.0 mg, 1 μmol) dissolved in DMSO (750 μL). The mixture was heated (55 °C, 25 min.) and remaining EDTA-anhydride was hydrolyzed (0.1M NaOH, 3mL, 55 °C, 10 min.). Aqueous TFA (0.1% wt/v) was added to adjust the total volume to 8 mL and the solution purified directly by reversed phase HPLC to provide 5-R as a white powder upon lyophilization of the appropriate fractions (0.6 mg, 60% recovery). MALDI-TOF-MS (monoisotopic), 1266.4 (1266.6 calc. for C<sub>57</sub>H<sub>76</sub>N<sub>19</sub>O<sub>15</sub>). The 5-R polyamide is shown in Figure 3.

**P. ImPyPy-(S)<sup>EDTA</sup>γ-PyPyPy-β-Dp (5-S)**

ImPyPy-(S)<sup>EDTA</sup>γ-PyPyPy-β-Dp was prepared from 1-S as described for 5-R (6.8 mg, 16% recovery). MALDI-TOF-MS (monoisotopic), 1266.5 (1266.6 calc. for C<sub>57</sub>H<sub>76</sub>N<sub>19</sub>O<sub>15</sub>).

**Example 2**

**Binding Site Size and Location by MPE•Fe(II) Footprinting**

**A. Preparation of 3'- and 5'-End-Labeled Restriction Fragments**

66074760  
15  
66074760  
The plasmid pMM5 was linearized with *Eco*RI and *Bsr*BI, then treated with the Sequenase enzyme, deoxyadenosine 5'-[ $\gamma$ - $^{32}$ P]triphosphate and thymidine 5'-[ $\gamma$ - $^{32}$ P]triphosphate for 3' labeling. Alternatively, pMM5 was linearized with *Eco*RI, treated with calf alkaline phosphatase, and then 5' labeled with T4 polynucleotide kinase and deoxyadenosine 5'-[ $\gamma$ - $^{32}$ P]triphosphate. The 5' labeled fragment was then digested with *Bsr*BI. The labeled fragment (3' or 5') was loaded onto a 6% non-denaturing polyacrylamide gel, and the desired 135 base pair band was visualized by autoradiography and isolated. Chemical sequencing reactions were performed according to published methods (Iverson, et al. *Nucl. Acids Res.* 1987, 15, 7823; Maxam, et al. *Methods Enzymol.* 1980, 65, 499).

#### B. MPE•Fe(II) Footprinting

15  
66074760  
All reactions were carried out in a volume of 40  $\mu$ L. A polyamide stock solution or water (for reference lanes) was added to an assay buffer where the final concentrations were: 25 mM Tris-acetate buffer (pH 7.0), 10 mM NaCl, 100  $\mu$ M/base pair calf thymus DNA, and 30 kcpm 3'- or 5'-radiolabeled DNA. The solutions were allowed to equilibrate for 4 hours. A fresh 50  $\mu$ M MPE•Fe(II) solution was prepared from 100  $\mu$ L of a 100  $\mu$ M MPE solution and 100  $\mu$ L of a 100  $\mu$ M ferrous ammonium sulfate ( $\text{Fe}(\text{NH}_4)_2(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$ ) solution. MPE•Fe(II) solution (5  $\mu$ M) was added to the equilibrated DNA, and the reactions were allowed to equilibrate for 5 minutes. Cleavage was initiated by the addition of dithiothreitol (5 mM) and allowed to proceed for 14 min. Reactions were stopped by ethanol precipitation, resuspended in 100 mM tris-borate-EDTA/80% formamide loading buffer, denatured at 85  $^{\circ}$ C for 6 min, and a 5  $\mu$ L sample (~ 15 kcpm) was immediately loaded onto an 8% denaturing polyacrylamide gel (5% crosslink, 7 M urea) at 2000 V.

#### C. Results

25  
MPE•Fe(II) footprinting (Van Dyke, et al. *Proc. Natl. Acad. Sci. U.S.A.* 1982, 79, 5470; Van Dyke, et al. *Science* 1984, 225, 1122) on 3'- and 5'- $^{32}$ P end-labeled 135 base

pair restriction fragments reveals that the polyamides, each at 1  $\mu$ M concentration, bind to the 5'-TGTTA-3' match site (25 mM Tris-acetate, 10 mM NaCl, 100  $\mu$ M/base pair calf thymus DNA, pH 7.0 and 22  $^{\circ}$ C) (Figure 5 and 6). Compounds **1-R** and **3-R** each at 1.25 M, protect both the cognate 5'-TGTA-3' site and the single base pair mismatch sequence 5'-TGTC-3'. Remarkably, binding sequence preferences vary for the polyamides depending on the stereochemistry of the amine substituent. At 1.25  $\mu$ M and 2.5 M concentration respectively, polyamides **1-S** and **3-S** bind a 5'-ACATT-3' reverse orientation match site in addition to the target match site 5'-TGTTA-3'. The sizes of the asymmetrically 3'-shifted footprint cleavage protection patterns for the polyamides are consistent with 5 base pair binding sites.

### **Example 3**

#### ***Binding Orientation Determination by Affinity Cleaving***

Affinity cleavage experiments (Taylor, et al. *Tetrahedron* **1984**, *40*, 457; Dervan, P. B. *Science* **1986**, *232*, 464) using hairpin polyamides modified with EDTA-Fe(II) at either the C-terminus or on the  $\gamma$ -turn, were used to determine polyamide binding orientation and stoichiometry. All reactions were carried out in a volume of 40  $\mu$ L. A polyamide stock solution or water (for reference lanes) was added to an assay buffer where the final concentrations were: 25 mM Tris-acetate buffer (pH 7.0), 20 mM NaCl, 100  $\mu$ M/base pair calf thymus DNA, and 20 kcpm 3'- or 5'-radiolabeled DNA. The solutions were allowed to equilibrate for 8 hours. A fresh solution of ferrous ammonium sulfate ( $\text{Fe}(\text{NH}_4)_2(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$ ) (10  $\mu$ M) was added to the equilibrated DNA, and the reactions were allowed to equilibrate for 15 minutes. Cleavage was initiated by the addition of dithiothreitol (10 mM) and allowed to proceed for 30 min. Reactions were stopped by ethanol precipitation, resuspended in 100 mM tris-borate-EDTA/80% formamide loading buffer, denatured at 85  $^{\circ}$ C for 6 min, and the entire sample was immediately loaded onto an 8% denaturing polyacrylamide gel (5% crosslink, 7 M urea) at 2000 V.

Affinity cleavage experiments were performed on the same 3'- and 5'-<sup>32</sup>P end-labeled 135 base pair restriction fragment (25 mM Tris-acetate, 10 mM NaCl, 100

μM/base pair calf thymus DNA, pH 7.0 and 22 °C). The observed cleavage patterns for

ImPyPy-(R)<sup>H2N</sup>γ-PyPyPy-β-Dp-EDTA•Fe(II) (4-R•Fe(II)), ImPyPy-

5 (R)EDTA•Fe(II) γ-PyPyPy-β-Dp (5-R•Fe(II)), ImPyPy-(S)<sup>H2N</sup>γ-PyPyPy-β-Dp-

EDTA•Fe(II) (4-S•Fe(II)), ImPyPy-(S)EDTA•Fe(II) γ-PyPyPy-β-Dp (5-S•Fe(II))

(Figures 7, 8 and 9) are in all cases 3'-shifted, consistent with minor groove occupancy.

In the presence of 3.3 μM of 4-R•Fe(II) and 10 μM 4-S•Fe(II) which have an

EDTA•Fe(II) moiety at the C-terminus, a single cleavage locus proximal to the 5' side of

the 5'-TGTTA-3' match sequence is revealed. In the presence of 3.3 μM 5-R•Fe(II) and

10 μM 5-S•Fe(II) which have an EDTA•Fe(II) moiety appended to the γ-turn, a single

cleavage locus is revealed proximal to the 3' side of the 5'-TGTTA-3' match sequence.

Cleavage loci are more concise for the γ-turn EDTA•Fe(II) placement relative to carboxy

terminal placement, consistent with the shorter tether. Cleavage loci are observed at both

the 5' and 3' side of the 5'-TGTTA-3' single base pair mismatch site in the presence of

10 μM of 4-R•Fe(II). The cleavage patterns observed at the 3' side of the site is

approximately 3-fold more intense than cleavage at the 5'-side. For polyamide 4-

S•Fe(II) at 10 μM concentration, a single cleavage locus is revealed proximal to the 5'

side of the 5'-ACATT-3' reverse orientation match site.

#### Example 4

#### *Energetics by Quantitative DNase I Footprinting*

##### A. DNase I Footprinting

All reactions were carried out in a volume of 400 μL. Carrier DNA was not used

in these reactions until after DNase I cleavage. A polyamide stock solution or water (for

reference lanes) was added to an assay buffer where the final concentrations were: 10

mM Tris•HCl buffer (pH 7.0), 10 mM KCl, 10 mM MgCl<sub>2</sub>, 5 mM CaCl<sub>2</sub>, and 30 kcpm

3'-radiolabeled DNA. The solutions were allowed to equilibrate for a minimum of 12

hours at 22 °C. Cleavage was initiated by the addition of 10 µL of a DNase I stock solution (diluted with 1 mM DTT to give a stock concentration of 1.875 u/mL) and was allowed to proceed for 7 min at 22 °C. The reactions were stopped by adding 50 µL of a solution containing 2.25 M NaCl, 150 mM EDTA, 0.6 mg/mL glycogen, and 30 µM base-pair calf thymus DNA, and then ethanol precipitated. The cleavage products were resuspended in 100 mM tris-borate-EDTA/80% formamide loading buffer, denatured at 85 °C for 6 min, and immediately loaded onto an 8% denaturing polyacrylamide gel (5% crosslink, 7 M urea) at 2000 V for 1 hour. The gels were dried under vacuum at 80 °C, then quantitated using storage phosphor technology.

Equilibrium association constants were determined as previously described (Mrksich, et al. *J. Am. Chem. Soc.* 1994, 116, 7983). The data were analyzed by performing volume integrations of the 5'-TGTTA-3' and 5'-TGACA-3 sites and a reference site. The apparent DNA target site saturation,  $\theta_{app}$ , was calculated for each concentration of polyamide using the following equation:

$$\theta_{app} = 1 - \frac{I_{tot}/I_{ref}}{I_{tot}^{\circ}/I_{ref}^{\circ}} \quad (1)$$

where  $I_{tot}$  and  $I_{ref}$  are the integrated volumes of the target and reference sites, respectively, and  $I_{tot}^{\circ}$  and  $I_{ref}^{\circ}$  correspond to those values for a DNase I control lane to which no polyamide has been added. The  $([L]_{tot}, \theta_{app})$  data points were fit to a Langmuir binding isotherm (eq 2,  $n=1$  for polyamides 1-3,  $n=2$  for polyamides 4 and 5) by minimizing the difference between  $\theta_{app}$  and  $\theta_{fit}$ , using the modified Hill equation:

$$\theta_{fit} = \theta_{min} + (\theta_{max} - \theta_{min}) \frac{K_a^n [L]_{tot}^n}{1 + K_a^n [L]_{tot}^n} \quad (2)$$

where  $[L]_{\text{tot}}$  corresponds to the total polyamide concentration,  $K_a$  corresponds to the equilibrium association constant, and  $\theta_{\text{min}}$  and  $\theta_{\text{max}}$  represent the experimentally determined site saturation values when the site is unoccupied or saturated, respectively. Data were fit using a nonlinear least-squares fitting procedure of KaleidaGraph software (version 2.1, Abelbeck software) with  $K_a$ ,  $\theta_{\text{max}}$ , and  $\theta_{\text{min}}$  as the adjustable parameters. All acceptable fits had a correlation coefficient of  $R > 0.97$ . At least three sets of acceptable data were used in determining each association constant. All lanes from each gel were used unless visual inspection revealed a data point to be obviously flawed relative to neighboring points. The data were normalized using the following equation:

$$\theta_{\text{norm}} = \frac{\theta_{\text{app}} - \theta_{\text{min}}}{\theta_{\text{max}} - \theta_{\text{min}}} \quad (3)$$

Photostimulable storage phosphorimaging plates (Kodak Storage Phosphor Screen S0230 obtained from Molecular Dynamics) were pressed flat against gel samples and exposed in the dark at 22 °C for 12-20 h. A Molecular Dynamics 400S PhosphorImager was used to obtain all data from the storage screens. The data were analyzed by performing volume integrations of all bands using the ImageQuant v. 3.2.

## B. Results

Quantitative DNase I footprint titrations (Brenowitz, et al., *Methods Enzymol.* 1986, 130, 132; Brenowitz, et al. *Proc. Natl. Acad. Sci. U.S.A.* 1986, 83, 8462; Senear, et al. *Biochemistry* 1986, 25, 7344) (10 mM Tris•HCl, 10 mM KCl, 10 mM MgCl<sub>2</sub> and 5 mM CaCl<sub>2</sub>, pH 7.0 and 22 °C) were performed to determine the equilibrium association constant ( $K_a$ ) of each six-ring hairpin polyamide for the three resolved sites (Figure 10 and 11). The 5'-TGTTA-3' site is bound by polyamides in the order: ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (1-R) ( $K_a = 3.8 \times 10^9 \text{ M}^{-1}$ ) > ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ -EtOH (2-R) ( $K_a = 3.3 \times 10^9 \text{ M}^{-1}$ ) > ImPyPy-(R)<sup>Ac</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (3-R) ( $K_a = 3.0 \times 10^8 \text{ M}^{-1}$ ) > ImPyPy- $\gamma$ -PyPyPy- $\beta$ -Dp ( $K_a = 2.9 \times 10^8 \text{ M}^{-1}$ ) > ImPyPy-(S)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (1-S) ( $K_a = 2.2 \times 10^7$

$M^{-1}) > \text{ImPyPy-(S)}^{\text{Ac}}\gamma\text{-PyPyPy}\beta\text{-Dp (3-S)} (K_a < 5.0 \times 10^6 M^{-1})$ . Equilibrium association constants for recognition of the 5'-TGACT-3' single base pair mismatch site are:  
 $\text{ImPyPy-(R)}^{\text{H}_2\text{N}}\gamma\text{-PyPyPy}\beta\text{-Dp (1-R)} (K_a = 3.5 \times 10^7 M^{-1})$   $_{-} \text{ImPyPy-(R)}^{\text{H}_2\text{N}}\gamma\text{-PyPyPy}\beta\text{-EtOH (2-R)} (K_a = 3.1 \times 10^7 M^{-1}) > \text{ImPyPy-(R)}^{\text{Ac}}\gamma\text{-PyPyPy}\beta\text{-Dp (3-R)} (K_a < 5 \times 10^6 M^{-1})$   
 $_{-} \text{ImPyPy-}\gamma\text{-PyPyPy}\beta\text{-Dp (K}_a = 4.8 \times 10^6 M^{-1})$ . The polyamides  $\text{ImPyPy-(S)}^{\text{H}_2\text{N}}\gamma\text{-PyPyPy}\beta\text{-Dp (1-S)}$  and  $\text{ImPyPy-(S)}^{\text{Ac}}\gamma\text{-PyPyPy}\beta\text{-Dp (3-S)}$  recognize the 5'-ACATT-3' reverse orientation sequence with  $K_a = 4.6 \times 10^6 M^{-1}$  and  $K_a < 5 \times 10^6 M^{-1}$  respectively. It should be noted that a detailed comparison of the relative mismatch binding energetics cannot be made since the 5'-TGACA-3' and 5'-ACATT-3' binding sites overlap. The relative affinity of 5'-TGTTA-3' match site binding varies from 100-fold to 5-fold depending on the stereochemistry of the  $\gamma$ -turn substitutions (Table 2).

**TABLE 2**  
*Equilibrium Association Constants ( $M^{-1}$ )<sup>a,b</sup>*

Improved Polyamide	Match Site 5'-TGTTA-3'	Reverse Site 5'-ACATT-3'	Mismatch Site 5'-TGACA-3'	Specificity <sup>c</sup>
ImPyPy- $\gamma$ -PyPyPy $\beta$ -Dp	$2.9 \times 10^8$	ND	$4.8 \times 10^6$	60
ImPyPy-(R) <sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy $\beta$ -Dp	$3 \times 10^9$ (0.2)	ND	$3.5 \times 10^7$ (1.0)	100
ImPyPy-(S) <sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy $\beta$ -Dp	$2.2 \times 10^7$ (0.7)	$4.6 \times 10^6$ (2.0) <sup>d</sup>	ND	5
ImPyPy-(R) <sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy $\beta$ -EtOH	$3.3 \times 10^9$ (0.9)	ND	$3.1 \times 10^7$ (0.4)	100
ImPyPy-(R) <sup>Ac</sup> $\gamma$ -PyPyPy $\beta$ -Dp	$3.0 \times 10^8$ (1.3)	ND	$< 5.0 \times 10^6$	$\geq 60$
ImPyPy-(S) <sup>Ac</sup> $\gamma$ -PyPyPy $\beta$ -Dp	$< 5.0 \times 10^6$	$< 5.0 \times 10^6$	ND	ND

<sup>a</sup>The reported association constants are the average values obtained from three DNase I footprint titration experiments. The standard deviation for each data set is indicated in parentheses. The assays were carried out at 22°C at pH 7.0 in the presence of 10 mM Tris-HCl, 10 mM KCl, 10 mM MgCl<sub>2</sub>, and 5 mM CaCl<sub>2</sub>.

<sup>b</sup>The five base pair binding sites are in capital letters.

<sup>c</sup>Specificity is calculated by  $K_a$  (match site)/ $K_a$  (mismatch site).

<sup>d</sup>Mismatch site is 5'(ACATT)-3' for ImPyPy-(S)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy $\beta$ -Dp<sub>(1-S)</sub> and ImPyPy-(S)<sup>Ac</sup> $\gamma$ -PyPyPy $\beta$ -Dp (3-S) as determined by MPE•Fe(II) footprinting and affinity cleaving.

ND = not determined.



### Example 5

#### *Binding site size and orientation*

MPE•Fe(II) footprinting reveals that the polyamides bind with highest affinity to the 5'-TGTTA-3' match site, the 5'-TGACA-3' single base pair mismatch site for polyamides 1-*R* and 3-*R*, and the 5'-ACATT-3' reverse orientation match site for polyamides 1-*S* and 3-*S* (Figure 6). Affinity cleaving experiments using polyamides with EDTA•Fe(II) placed at either the carboxy terminus or the  $\gamma$ -turn confirm that polyamides derived from both (*R*) and (*S*)-2,4-diaminobutyric acid bind to the 5'-TGTTA-3' target site with a single orientation (Figure 10). The observation of a single cleavage locus is consistent only with an oriented 1:1 polyamide:DNA complex in the minor groove and rules out any dimeric overlapped or extended binding motifs. The hairpin binding model is further supported by the location of the cleavage locus at either the 5' or 3' side of the 5'-TGTTA-3' target site corresponding to EDTA•Fe(II) placement at the polyamide carboxy terminus or the  $\gamma$ -turn, respectively (Figure 10). Polyamide subunits linked by the (*R*)<sup>H<sub>2</sub>N</sup> $\gamma\gamma$  bind the symmetric single base pair mismatch sequence 5'-TGACA-3' in two distinct orientations. Polyamides linked with (*S*)<sup>H<sub>2</sub>N</sup> $\gamma$  bind to a 5'-ACATT-3' reverse orientation match sequence as revealed by a unique cleavage locus at the 5' side of the site.

### Example 6

#### *Binding Affinity*

All six polyamides bind to the 5'-TGTTA-3' target site with binding isotherms (eq. 2,  $n = 1$ ) consistent with binding as an intramolecular hairpin (Figure 11). However the relative match site binding affinity varies by nearly 1000-fold depending on the stereochemistry of the  $\gamma$ -turn and the nature of the substituents. Among the six polyamides, ImPyPy-(*R*)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (1-*R*) binds to the targeted 5'-TGTTA-3' site with the highest affinity. ImPyPy-(*R*)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ -Dp binds with an equilibrium association constant, ( $K_a = 3 \times 10^9 \text{ M}^{-1}$ ; Parks, et al. *J. Am. Chem. Soc.* 1996, 118, 6147),

a factor of 10 greater than that of the parent polyamide, ImPyPy- $\gamma$ -PyPyPy- $\beta$ -Dp, ( $K_a = 3 \times 10^8 \text{ M}^{-1}$ ). Replacement of the C-terminal dimethylaminopropylamide group of **1-R** with an ethoxyamide group as in ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ -EtOH (**2-R**) results in no decrease in binding affinity ( $K_a = 3 \times 10^9 \text{ M}^{-1}$ ). Acetylation of the  $\gamma$ -turn amino group as  
 5 in ImPyPy-(R)<sup>Ac</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (**3-R**) reduces binding affinity 10-fold ( $K_a = 3 \times 10^8 \text{ M}^{-1}$ ) relative to **1-R**.

The observation that polyamides which differ only by replacement of the dimethylaminopropylamide group **1-R** with an ethoxyamide group **2-R** bind with similar affinity indicates that interactions between the cationic dimethylaminopropyl tail group with anionic phosphate residues or the negative electrostatic potential in the floor of the minor groove may not contribute substantially to the energetics of hairpin-DNA binding (Zimmer, et al. *Prog. Biophys. Molec. Biol.* **1986**, *47*, 31; Pullman, B. *Adv. Drug. Res.* **1990**, *18*, 1; Breslauer, et al. *Structure and Expression (Vol. 2), DNA and Its Drug Complexes* p. 273-289, R. H. Sarma and M. H. Sarma (eds.) Academic Press (1988)).  
 10 Furthermore, these results indicate that the observed binding enhancement of **1-R**, in relation to ImPyPy- $\gamma$ -PyPyPy- $\beta$ -Dp, is not simply the difference between a monocationic and dicationic ligand binding to the polycationic DNA helix (Zimmer, et al. *Prog. Biophys. Molec. Biol.* **1986**, *47*, 31; Pullman, B. *Adv. Drug. Res.* **1990**, *18*, 1; Breslauer, et al. *Structure and Expression (Vol. 2), DNA and Its Drug Complexes* p. 273-289, R. H.  
 15 Sarma and M. H. Sarma (eds.) Academic Press (1988)). The modest increased binding affinity of polyamide **1-R** may result from electrostatic interactions between the precisely placed amine group and the floor of the minor groove. Alternately, the increased affinity could indicate a reduction in the degrees of freedom accessible to the free hairpin in solution resulting from a steric effect, or an electrostatic interaction between the  
 20 positively charged amine group and the negative potential of the  $\gamma$ -carbonyl group.

Polyamides linked with (S)<sup>H<sub>2</sub>N</sup> $\gamma$ , ImPyPy-(S)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (**1-S**) and ImPyPy-(S)<sup>Ac</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (**3-S**), bind to the 5'-TGTTA-3' match site with 100-fold

( $K_a = 2 \times 10^7 \text{ M}^{-1}$ ) and 1000-fold ( $K_a < 5 \times 10^6 \text{ M}^{-1}$ ) reduced affinity relative to the ( $R$ )<sup>H<sub>2</sub>N</sup>γ linked polyamide **1-R**. These results demonstrate that the DNA-binding affinity of chiral hairpin polyamides can be predictably regulated as a function of the stereochemistry of the turn residue.

#### Example 7

##### *Sequence-specificity*

Polyamides with a variety of substitutions at the γ-turn bind preferentially to the 5'-TGTTA-3' match site, while overall specificity versus binding at reverse orientation and mismatch sites is modified. Replacing the α-proton in the γ-residue of ImPyPy-γ-PyPyPy-β-Dp with an amino group that confers a chiral α-hydrogen ( $R$ ) configuration, provides the most specific polyamide ImPyPy-( $R$ )<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp (**1-R**). The ImPyPy-( $R$ )<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp•5'-TGTTA-3' complex forms with 100-fold preference relative to the ImPyPy-( $R$ )<sup>H<sub>2</sub>N</sup>γ-PyPyPy-β-Dp•5'-TGTC A-3' mismatch complex. Substitution of the charged dimethylaminopropyl tail group with an ethoxyamide group as in (**2-R**) does not alter binding specificity. The modest increase in specificity against single base mismatch sequences for polyamides **1-R** and **2-R** (100-fold) relative to the parent unsubstituted hairpin polyamide (60-fold) implicates chiral hairpin polyamides as an optimized class of small molecules for recognition of the DNA minor groove.

#### Example 8

##### *Binding Orientation*

In principle, a polyamide:DNA complex can form at two different DNA sequences depending on the alignment of the polyamide (N-C) with the walls of the minor groove (5'-3'; White, et al. *J. Am. Chem. Soc.* **1997**, *119*, 8756). A six ring-hairpin polyamide of core sequence composition ImPyPy-γ-PyPyPy which places the N-terminus of each three-ring polyamide subunit at the 5'-side of individual recognized DNA strands would bind to 'forward match' 5'-WGWWW-3' sequences (W = A or T). Placement of the polyamide N-terminus at the 3'-side of each recognized strand would result in

targeting 'reverse match' 5'-WCWW-3' sequences. For hairpin polyamides there is an energetic preference for 'forward' alignment of each polyamide subunit (N-C) with respect to the backbone (5'-3') of the DNA double helix (White, et al. *J. Am. Chem. Soc.* 1997, 119, 8756).

5 In addition to decreasing the affinity for the 5'-TGTTA-3' match site, replacing the  $\alpha$ -proton of  $\gamma$ -turn in ImPyPy- $\gamma$ -PyPyPy- $\beta$ -Dp with (*S*)<sup>H<sub>2</sub>N</sup> $\gamma$  changes the mismatch sequence preference from the 5'-TGACA-3' site bound by the (*R*)<sup>H<sub>2</sub>N</sup> $\gamma$ -linked polyamides to a 5'-ACATT-3' reverse match site. Binding to the reverse site may result from the presence of the steric bulk of the amino or acetamido groups in the floor of the minor groove preventing the deep polyamide binding required for specific DNA recognition. 10 However, an analysis of hairpin folding requirements for 'forward' and 'reverse' binding reveals an additional model.

In principle, there exist two non-superimposable hairpin folds which are related by mirror plane symmetry (Figure 12). One hairpin fold is responsible for the preferred 15 5' to 3' N to C orientation, while the other fold corresponds to the 3' to 5' N to C reverse orientation binding. For an achiral hairpin polyamide in the absence of DNA, each non-superimposable fold should be energetically equivalent. However, an asymmetrically folded hairpin polyamide with a chiral substituent could potentially display differential energetics for oriented binding (Figure 12). In the forward folded hairpin (5' to 3' N to 20 C), (*R*)<sup>H<sub>2</sub>N</sup> $\gamma$  directs the amine functionality away from the DNA helix, while (*S*) enantiomer is predicted to direct the amine into the floor of the minor groove. For the 'reverse' fold hairpin, (*S*)<sup>H<sub>2</sub>N</sup> $\gamma$  directs the amine functionality away from the floor of the DNA helix, while the amine of the (*R*) enantiomer is predicted to clash with the floor of the helix. The modest enhanced specificity of chiral polyamides **1-R** and **2-R** relative to 25 the unsubstituted parent hairpin may result from stabilization of the forward binding mode and/or destabilization of the reverse binding hairpin fold.

## Example 9

### Tandemly-Linked Polyamides

#### A. Synthesis

5 ImPyPy-(R)[ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (6) and ImPyPy-(R)[ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\delta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (7) were synthesized from Boc- $\beta$ -alanine-Pam resin (0.6 g resin, 0.6 mmol/g substitution) using Boc-chemistry machine-assisted protocols in 31 steps (Figure 3).<sup>[12]</sup> ImPyPy-(R)<sup>FmocHN</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-Pam resin was prepared as described.<sup>[10]</sup> The Fmoc protecting group was then removed by treatment with (4:1) piperidine/DMF. The remaining amino acid sequence was then synthesized in a stepwise manner using Boc-chemistry machine assisted protocols to provide ImPyPy-(R)[ImPyPy-(R)<sup>FmocHN</sup> $\gamma$ -PyPyPy- $\beta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-Resin and ImPyPy-(R)[ImPyPy-(R)<sup>FmocHN</sup> $\gamma$ -PyPyPy- $\delta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-Resin. The Fmoc group was removed with (4:1) piperidine/DMF. A sample of resin was then cleaved by a single-step aminolysis reaction with ((dimethylamino)propylamine (55 °C, 18 h) and the reaction mixture subsequently purified by reversed phase HPLC to provide ImPyPy-(R)[ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (6) and ImPyPy-(R)[ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\delta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (7). For the synthesis of the EDTA-turn derivative 7-E, a sample ImPyPy-(R)[ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\delta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (7) was treated with an excess of EDTA-dianhydride (DMSO/NMP, DIEA 55 °C, 30 min.) and the remaining anhydride hydrolyzed (0.1 M NaOH, 55 °C, 10 min.). The polyamide ImPyPy-(R)[ImPyPy-(R)<sup>EDTA</sup> $\gamma$ -PyPyPy- $\delta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (7-E) was then isolated by reverse phase HPLC. The dicationic twelve-ring tandem hairpins are soluble at concentrations up to 1 mM. The solubility of the tandem hairpins is 10- to 100-fold greater than that found for extended or hairpin twelve-ring polyamides.

#### ImPyPy-(R)[ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Dp ("6")

30 ImPyPy-(R)<sup>Fmoc</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-Resin was synthesized in a stepwise fashion by machine-assisted solid phase methods from Boc- $\beta$ -Pam-Resin (0.6 mmol/g). (R)-2-Fmoc-4-Boc-diaminobutyric acid (0.7 mmol) was incorporated as previously

described for Boc- $\gamma$ -aminobutyric acid. ImPyPy-(R)<sup>Fmoc</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-Resin was placed in a glass 20 mL peptide synthesis vessel and treated with DMF (2 mL), followed by piperidine (8 mL) and agitated (22 °C, 30 min.). ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-resin was isolated by filtration, and washed sequentially with an excess of DMF, DCM, MeOH, and ethyl ether and the amine-resin dried *in vacuo*. ImPyPy-(R)[ImPyPy-(R)<sup>FmocHN</sup> $\gamma$ -PyPyPy- $\beta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-Resin was then synthesized in a stepwise fashion by machine-assisted solid phase methods from ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-resin (0.38 mmol/g). ImPyPy-(R)[ImPyPy-(R)<sup>FmocHN</sup> $\gamma$ -PyPyPy- $\beta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-Resin was placed in a glass 20 mL peptide synthesis vessel and treated with DMF (2 mL), followed by piperidine (8 mL) and agitated (22 °C, 30 min.). ImPyPy-(R)[ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-Resin was isolated by filtration, and washed sequentially with an excess of DMF, DCM, MeOH, and ethyl ether and the amine-resin dried *in vacuo*. A sample of ImPyPy-(R)[ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-Resin (240 mg, 0.29 mmol/gram) was treated with neat dimethylaminopropylamine (2 mL) and heated (55 °C) with periodic agitation for 16 h. The reaction mixture was then filtered to remove resin, 0.1% (wt/v) TFA added (6 mL) and the resulting solution purified by reversed phase HPLC. ImPyPy-(R)[ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Dp is recovered upon lyophilization of the appropriate fractions as a white powder (28 mg, 22% recovery). UV (H<sub>2</sub>O)  $\lambda_{\text{max}}$  246, 306 (100,000); MALDI-TOF-MS [M<sup>+</sup>-H] (monoisotopic), 1881.9: 1881.9 calc. for C<sub>89</sub>H<sub>109</sub>N<sub>32</sub>O<sub>16</sub>

**ImPyPy-(R)[ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\delta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Dp ("7")**

ImPyPy-(R)[ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\delta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-Resin was prepared as described for ImPyPy-(R)[ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-Resin. A sample of ImPyPy-(R)[ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\delta$ ]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Pam-Resin (240 mg, 0.29 mmol/gram<sup>19</sup>) was treated with neat dimethylaminopropylamine (2 mL) and heated (55 °C) with periodic agitation for 16 h. The reaction mixture was then filtered to remove resin, 0.1% (wt/v) TFA added (6 mL) and the resulting solution purified by reversed phase HPLC. ImPyPy-

(R)[ImPyPy-(R)<sup>H2N</sup> $\gamma$ -PyPyPy- $\delta$ -]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Dp is recovered upon lyophilization of the appropriate fractions as a white powder (32 mg, 25% recovery).  $[\alpha]_D^{20} +14.6$  (c 0.05, H<sub>2</sub>O); UV (H<sub>2</sub>O)  $\lambda_{max}$  246, 306 (100,000); <sup>1</sup>H NMR (300 MHz, [D<sub>6</sub>]DMSO, 20 °C):  $\delta$  = 10.54 (s, 1 H; aromatic NH); 10.45 (s, 1 H; aromatic NH); 10.44 (s, 1 H; aromatic NH); 10.02 (s, 1 H; aromatic NH); 9.95 (s, 1 H; aromatic NH); 9.92 (s, 1 H; aromatic NH); 9.90 (d, 2 H; aromatic NH); 9.86 (d, 2 H; aromatic NH); 9.2 (br s, 1 H; CF<sub>3</sub>COOH); 8.25 (m, 4 H; aliphatic NH, NH<sub>3</sub>); 8.11 (d, 1 H; *J* = 8.5 Hz, aliphatic NH); 8.04 (m, 4H, aliphatic NH), 7.37 (s, 2 H; CH); 7.25 (m, 2 H; CH); 7.22 (d, 1 H; CH); 7.18 (m, 2 H; CH); 7.16 (m, 3 H; CH); 7.12 (m, 4 H; CH); 7.02 (m, 4 H; CH); 6.95 (d, 1 H; *J* = 1.6 Hz; CH); 6.91 (d, 1 H; *J* = 1.5 Hz; CH); 6.88 (d, 1 H, *J* = 1.3 Hz; CH); 6.85 (m, 3 H; CH); 5.32 (t, 1 H; aliphatic CH), 4.45 (m, 1 H, aliphatic CH), 3.96 (s, 6 H; NCH<sub>3</sub>); 3.83 (s, 3 H; NCH<sub>3</sub>); 3.80 (s, 18 H; NCH<sub>3</sub>); 3.79 (s, 3 H; NCH<sub>3</sub>); 3.76 (s, 3 H; NCH<sub>3</sub>); 3.39 (m, 4 H; CH<sub>2</sub>); 3.28 (m, 2 H; CH<sub>2</sub>); 3.15 (m, 4 H; CH<sub>2</sub>); 3.07 (m, 2 H; CH<sub>2</sub>); 2.97 (m, 2 H; CH<sub>2</sub>); 2.70 (d, 6 H; N(CH<sub>3</sub>)<sub>2</sub>); 2.32 (m, 2 H; CH<sub>2</sub>); 1.93 (m, 2 H; CH<sub>2</sub>); 1.71 (m, 2 H; CH<sub>2</sub>); 1.47 (m, 2 H; CH<sub>2</sub>); 1.20 (m, 4 H; CH<sub>2</sub>); MALDI-TOF-MS [*M*<sup>+</sup>-H] (monoisotopic), 1910.2: 1909.9 calc. for C<sub>91</sub>H<sub>113</sub>N<sub>32</sub>O<sub>16</sub>.

**ImPyPy-(R)[ImPyPy-(R)<sup>EDTA</sup> $\gamma$ -PyPyPy- $\delta$ -]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Dp ("7-E")**

Excess EDTA-dianhydride (50 mg) was dissolved in DMSO/NMP (1 mL) and DIEA (1 mL) by heating at 55 °C for 5 min. The dianhydride solution was added to ImPyPy-(R)[ImPyPy-(R)<sup>H2N</sup> $\gamma$ -PyPyPy- $\delta$ -]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (10 mg, 5  $\mu$ mol) dissolved in DMSO (750  $\mu$ L). The mixture was heated (55 °C, 25 min.) and the remaining EDTA-anhydride hydrolyzed (0.1M NaOH, 3 mL, 55 °C, 10 min). Aqueous TFA (0.1% wt/v) was added to adjust the total volume to 8 mL and the solution purified directly by reversed phase HPLC to provide ImPyPy-(R)[ImPyPy-(R)<sup>EDTA</sup> $\gamma$ -PyPyPy- $\delta$ -]<sup>HN</sup> $\gamma$ -PyPyPy- $\beta$ -Dp (7-E) as a white powder upon lyophilization of the appropriate fractions (2 mg, 20% recovery). MALDI-TOF-MS [*M*<sup>+</sup>-H] (monoisotopic), 2184.3: 2184.0 calc. for C<sub>101</sub>H<sub>127</sub>N<sub>34</sub>O<sub>23</sub>

## B. Plasmid and restriction fragment preparation

The plasmids pDH10, pDH11, and pDH12 were constructed by hybridization of the inserts listed in Figure 18. Each hybridized insert was ligated individually into linearized pUC19 *Bam*HI/*Hind*III plasmid using T4 DNA ligase. The resultant constructs were used to transform Top10F' OneShot competent cells from Invitrogen. Ampicillin-resistant white colonies were selected from 25 mL Luria-Bertani medium agar plates containing 50 µg/mL ampicillin and treated with XGAL and IPTG solutions. Large-scale plasmid purification was performed with Qiagen Maxi purification kits. Dideoxy sequencing was used to verify the presence of the desired insert. Concentration of the prepared plasmid was determined at 260 nm using the relationship of 1 OD unit = 50 µg/mL duplex DNA.

The plasmids pDH(11-12) were linearized with *Eco*RI and *Bsr*BI, then treated with the Sequenase enzyme, deoxyadenosine 5'-[α-<sup>32</sup>P]triphosphate and thymidine 5'-[α-<sup>32</sup>P]triphosphate for 3' labeling. Alternatively, these plasmids were linearized with *Eco*RI, treated with calf alkaline phosphatase, and then 5' labeled with T4 polynucleotide kinase and deoxyadenosine 5'-[γ-<sup>32</sup>P]triphosphate. The 5' labeled fragment was then digested with *Bsr*BI. The labeled fragment (3' or 5') was loaded onto a 6% non-denaturing polyacrylamide gel, and the desired 147 base pair band was visualized by autoradiography and isolated. Chemical sequencing reactions were performed according to published methods.

## C. Binding Site Size

All reactions were carried out in a volume of 40 µL. A polyamide stock solution or water (for reference lanes) was added to an assay buffer where the final concentrations were: 25 mM Tris-acetate buffer (pH 7.0), 10 mM NaCl, 100 µM/base pair calf thymus DNA, and 30 kcpm 3'- or 5'-radiolabeled DNA. The solutions were allowed to equilibrate for 4 hours. A fresh 50 µM MPE•Fe(II) solution was prepared from 100 µL of a 100 µM MPE solution and 100 µL of a 100 µM ferrous ammonium sulfate (Fe(NH<sub>4</sub>)<sub>2</sub>(SO<sub>4</sub>)<sub>2</sub>•6H<sub>2</sub>O) solution. MPE•Fe(II)



00374702 081209 061800 20242000

5 solution (5  $\mu$ M) was added to the equilibrated DNA, and the reactions were allowed to equilibrate for 5 minutes. Cleavage was initiated by the addition of dithiothreitol (5 mM) and allowed to proceed for 14 min. Reactions were stopped by ethanol precipitation, resuspended in 100 mM tris-borate-EDTA/80% formamide loading buffer, denatured at 85 °C for 6 min, and a 5  $\mu$ L sample (~ 15 kcpm) was immediately loaded onto an 8% denaturing polyacrylamide gel (5% crosslink, 7 M urea) at 2000 V.

10 MPE•Fe(II) footprinting on 3'- or 5'-<sup>32</sup>P end-labeled 135 base pair *Eco*RI/*Bsr*BI restriction fragments from the plasmid pDH11 reveals that polyamide 7, at 100 pM concentration, binds to the designated 11-bp match site 5'-TGTTATTGTTA-3' (25 mM Tris-acetate, 10 mM NaCl, pH 7.0 and 22 °C) (Figures 5a and 5c). Binding of the mismatch site 5'-TGTCATTGTCA-3' is only observed at much higher polyamide concentrations (Figure 5a). The size of the asymmetrically 3'-shifted cleavage protection pattern for polyamide 7 at the designated match site 5'-TGTTATTGTTA-3' is consistent with formation of the predicted hairpin- $\delta$ -hairpin•DNA complex.

#### D. Binding Orientation

20 All reactions were carried out in a volume of 40  $\mu$ L. A polyamide stock solution or water (for reference lanes) was added to an assay buffer where the final concentrations were: 25 mM Tris-acetate buffer (pH 7.0), 20 mM NaCl, 100  $\mu$ M/base pair calf thymus DNA, and 20 kcpm 3'- or 5'-radiolabeled DNA. The solutions were allowed to equilibrate for 8 hours. A fresh solution of ferrous ammonium sulfate ( $\text{Fe}(\text{NH}_4)_2(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$ ) (10  $\mu$ M) was added to the equilibrated DNA, and the reactions were allowed to equilibrate for 15 minutes. Cleavage was initiated by the addition of

25 dithiothreitol (10 mM) and allowed to proceed for 30 min. Reactions were stopped by ethanol precipitation, resuspended in 100 mM tris-borate-EDTA/80% formamide loading buffer, denatured at 85 °C for 6 min, and the entire sample was immediately loaded onto an 8% denaturing polyacrylamide gel (5% crosslink, 7 M urea) at 2000 V.

30 Affinity cleavage experiments using 7-E which has an EDTA•Fe(II) moiety appended to the  $\gamma$ -turn, were used to confirm polyamide binding orientation and

stoichiometry. Affinity cleavage experiments were performed on the same 3'- or 5'-<sup>32</sup>P end-labeled 135 base pair DNA restriction fragment from the plasmid pDH11 (25 mM Tris-acetate, 10 mM NaCl, 100  $\mu$ M/base pair calf thymus DNA, pH 7.0 and 22 °C). The observed cleavage pattern for 7-E (Figures 5b and 5d) are 3'-shifted, consistent with minor groove occupancy. In the presence of 1  $\mu$ M 7-E, a single cleavage locus proximal to the 3' side of the 5'-TGTTATTGTTA-3' match sequence is revealed, consistent with formation of an oriented 1:1 hairpin- $\delta$ -hairpin•DNA complex.







#### 10 E. Equilibrium Association Constants

Quantitative DNase I footprinting and related mathematical calculations were performed as described above in Example 4, except as otherwise indicated below. Quantitative DNase I footprint titrations (10 mM Tris•HCl, 10 mM KCl, 10 mM MgCl<sub>2</sub> and 5 mM CaCl<sub>2</sub>, pH 7.0 and 22 °C) were performed to determine the equilibrium association constants ( $K_a$ ) of 6 and 7 for the 10-, 11- and 12-bp match and mismatch sites (Table 1). Polyamide 7 preferentially binds the 11-bp 5'-TGTTATTGTTA-3' target sequence with an equilibrium association constant,  $K_a > 1 \times 10^{11} \text{ M}^{-1}$ . The corresponding 11 bp mismatch 5'-TGTCATTGTCA-3' site is bound by 7 with  $> 4500$ -fold lower affinity ( $K_a = 2.2 \times 10^8 \text{ M}^{-1}$ ). Polyamide 7 binds the 10 bp site 5'-TGTTATTGTTA-3' ( $K_a = 1.5 \times 10^{10} \text{ M}^{-1}$ ) and the 12 bp site 5'-TGTTATATGTTA-3' ( $K_a = 1.0 \times 10^9 \text{ M}^{-1}$ ) with 70- and 1000-fold lower affinity, respectively. Polyamide 6 binds the 10-bp 5'-TGTTATTGTTA-3' site and 11-bp 5'-TGTTATTGTTA-3' site with  $K_a = 2 \times 10^{10} \text{ M}^{-1}$ , and also binds the 12-bp 5'-TGTTATATGTTA-3' site with 16-fold lower affinity ( $K_a = 9.0 \times 10^9 \text{ M}^{-1}$ ). The parent hairpin ImPyPy-(R)<sup>H<sub>2</sub>N</sup> $\gamma$ -PyPyPy- $\beta$ -Dp was found to bind to the 5'-TGTTA-3' match site with  $K_a = 5 \times 10^9 \text{ M}^{-1}$ .

Formally the subunits of polyamide 7 represent the combination of the parent-acetylated parent tandem hairpin. The parent and acetylated hairpins occupy I with binding energetics of -13.2 kcal/mol and -11.8 kcal/mol respectively, predicting that covalent linkage of parent- $\delta$ -acetylated would bind the same site

with an association constant of ( $K_a$ ) =  $2.2 \times 10^{18} \text{ M}^{-1}$ , 6 orders of magnitude higher than the observed tandem hairpin 7 which binds with -16.3 kcal/mol. Relative to parent recognition of the 5'-TGTTA-3' half site, we only observe a 2-fold enhancement and 5-fold decrease in binding respectively, recognition of the 10- and 12-bp sites. At site IV, a single base pair mismatch reduces binding energetics for both unlinked and linked hairpins. Unlinked parent and acetylated-parent respectively bind with 10.5 kcal/mol and 9.2 kcal/mol, predicting the linked polyamide would bind with a ( $K_a$ ) =  $2.4 \times 10^{14} \text{ M}^{-1}$ . The observed tandem hairpin binds with energetics of -11.35 kcal/mol and a ( $K_a$ ) =  $2.2 \times 10^8 \text{ M}^{-1}$ .

**Table 2** Equilibrium Association Constants ( $\text{M}^{-1}$ )<sup>a,b</sup>

Polyamide	5'-aTGTTATGTTAg-3'	5'-aTGTCATGTCA-3'	Specificity
	<b><math>2 \times 10^{10}</math></b>	$1.5 \times 10^8$	133
	<b><math>1.5 \times 10^{10}</math></b>	$1.9 \times 10^8$	80
Polyamide	5'-aTGTTATTGTTAg-3'	5'-aTGTCATTGTCA-3'	Specificity
	<b><math>1.5 \times 10^{10}</math></b>	$2 \times 10^8$	75
	<b><math>\geq 1 \times 10^{12}</math></b>	$2.2 \times 10^8$	$\geq 4500$
Polyamide	5'-aTGTTATATGTTAg-3'	5'-aTGTCATATGTCA-3'	Specificity
	<b><math>9 \times 10^8</math></b>	$3 \times 10^7$	30
	<b><math>1 \times 10^9</math></b>	$2.5 \times 10^7$	40

<sup>a</sup>The reported association constants are the average values obtained from three DNase I footprint titration experiments. The assays were carried out at 22 °C at pH 7.0 in the presence of 10 mM Tris-HCL, 10 mM KCl, 10 mM MgCl<sub>2</sub>, and 5 mM CaCl<sub>2</sub>. <sup>b</sup>The ten, eleven, and twelve base-pair sites are in capital letters, with flanking sequences in lower case letters. Match site association constants are shown in boldtype. Specificity is calculated as  $K_a(\text{match})/K_a(\text{mismatch})$ .

## F. Linker Dependence

Site size preferences of polyamides 6 and 7 result from modifications to the length of the turn-to-tail linker. Modeling indicated that  $\beta$  and  $\delta$  linkers would provide sufficient length for recognition of either 10- or 11- base pairs, but would be too short to span the 12-bp binding site. Polyamide 7 displays optimal recognition of the 11-bp site, binding the 5'-TGTTATTGTTA-3' site with a  $K_a = 1 \times 10^{12} \text{ M}^{-1}$ .

Replacing the  $\delta$  linker in 7 with the 2-carbon shorter  $\beta$ -alanine residue in 6 shows a reduction of affinity at the 11-bp site by > 6-fold ( $K_a = 1.5 \times 10^{10} \text{ M}^{-1}$ ). The highly reduced affinities of 6 and 7 at the 12-bp site indicates that covalent constraints of the linker subunit prevents alignment of hairpin subunits for their binding sites.

**Table 3**

*Binding affinity of 6-ring hairpin and  $\delta$ -linked tandem hairpins at 11-bp match site I, and a series of mismatch sites II-VI*

	Parent	Polyamide 2
I	<p><math>K_a = 5 \times 10^9 \text{ M}^{-1}</math></p>	<p><math>K_a = 1 \times 10^{12} \text{ M}^{-1}</math></p>
II	<p><math>K_a = 5 \times 10^9 \text{ M}^{-1}</math></p>	<p><math>K_a = 1.5 \times 10^{10} \text{ M}^{-1}</math></p>
III	<p><math>K_a = 5 \times 10^9 \text{ M}^{-1}</math></p>	<p><math>K_a = 1 \times 10^9 \text{ M}^{-1}</math></p>
IV	<p><math>K_a = 5 \times 10^7 \text{ M}^{-1}</math></p>	<p><math>K_a = 2.2 \times 10^8 \text{ M}^{-1}</math></p>
V	<p><math>K_a = 3 \times 10^8 \text{ M}^{-1}</math></p>	<p><math>K_a = 1 \times 10^8 \text{ M}^{-1}</math></p>
VI	<p><math>K_a = 1 \times 10^8 \text{ M}^{-1}</math></p>	<p><math>K_a = 5 \times 10^7 \text{ M}^{-1}</math></p>

As disclosed herein, the present invention provides the reagents and methodologies for the preparation and use of a variety of new chiral hairpin polyamide structures for specific recognition in the DNA minor groove. While a preferred form of the invention has been shown in the drawings and described, since variations in the preferred form will be apparent to those skilled in the art, the invention should not be construed as limited to the specific form shown and described, but instead is as set forth in the claims.

662782 2024260